



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C07K 14/00	A2	(11) International Publication Number: WO 00/23464 (43) International Publication Date: 27 April 2000 (27.04.2000)
(21) International Application Number: PCT/EP99/07742 (22) International Filing Date: 14 October 1999 (14.10.1999) (30) Priority Data: 09/173,941 16 October 1998 (16.10.1998) US (60) Parent Application or Grant NOVARTIS AG [/]; (). NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [/]; (). THE SCRIPPS RESEARCH INSTITUTE [/]; (). BARBAS, Carlos, F. [/]; (). BARBAS, Carlos, F. [/]; (). BECKER, Konrad ; ().	Published	

(54) Title: ZINC FINGER BINDING DOMAINS FOR GNN
(54) Titre: DOMAINES DE LIAISON DES DOIGTS DE ZINC POUR GNN

(57) Abstract

Zinc finger-nucleotide binding polypeptides having binding specificity for target nucleotides containing one or GNN triplets are provided. Compositions containing such polypeptides and the use of such polypeptides and compositions for regulating nucleotide function are also provided.

(57) Abrégé

L'invention porte sur des polypeptides de liaison des doigts de zinc ayant une spécificité de liaison pour des nucléotides cibles contenant un ou des triplets de GNN. L'invention porte également sur des compositions contenant ces polypeptides et sur l'utilisation de ces polypeptides et de ces compositions pour réguler la fonction nucléotidique.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/00	A2	(11) International Publication Number: WO 00/23464 (43) International Publication Date: 27 April 2000 (27.04.00)
(21) International Application Number: PCT/EP99/07742 (22) International Filing Date: 14 October 1999 (14.10.99) (30) Priority Data: 09/173,941 16 October 1998 (16.10.98) US (71) Applicant (for all designated States except AT US): NOVAR-TIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VER-WALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (71) Applicant (for all designated States except US): THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, La Jolla, CA 92037 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): BARBAS, Carlos, F. [US/US]; 755 Pacific Surf Drive, Solana Beach, CA 92075 (US).	(74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SI, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: ZINC FINGER BINDING DOMAINS FOR GNN (57) Abstract <p>Zinc finger-nucleotide binding polypeptides having binding specificity for target nucleotides containing one or GNN triplets are provided. Compositions containing such polypeptides and the use of such polypeptides and compositions for regulating nucleotide function are also provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5

10

15

20

25

30

35

40

45

50

55

5

ZINC FINGER BINDING DOMAINS FOR GNN

10

Technical Field of the Invention

15

The field of this invention is zinc finger protein binding to target nucleotides. More particularly, the present invention pertains to amino acid residue sequences within the α -helical domain of zinc fingers that specifically bind to target nucleotides of the formula 5'-(GNN)-3'.

20

Background of the Invention

25

30

35

40

45

50

55

The paradigm that the primary mechanism for governing the expression of genes involves protein switches that bind DNA in a sequence specific manner was established in 1967 (Ptashne, M. (1967) *Nature (London)* 214, 323-4). Diverse structural families of DNA binding proteins have been described. Despite a wealth of structural diversity, the Cys₂-His₂ zinc finger motif constitutes the most frequently utilized nucleic acid binding motif in eukaryotes. This observation is as true for yeast as it is for man. The Cys₂-His₂ zinc finger motif, identified first in the DNA and RNA binding transcription factor TFIIIA (Miller, J., McLachlan, A. D. & Klug, A. (1985) *Embo J* 4, 1609-14), is perhaps the ideal structural scaffold on which a sequence specific protein might be constructed. A single zinc finger domain consists of approximately 30 amino acids with a simple $\beta\beta\alpha$ fold stabilized by hydrophobic interactions and the chelation of a single zinc ion (Miller, J., McLachlan, A. D. & Klug, A. (1985) *Embo J* 4, 1609-14, Lee, M. S., Gippert, G. P., Soman, K. V., Case, D. A. & Wright, P. E. (1989) *Science* 245, 635-7). Presentation of the α -helix of this domain into the major groove of DNA allows for sequence specific base contacts. Each zinc finger domain typically recognizes three base pairs of DNA (Pavletich, N. P. & Pabo, C. O. (1991) *Science (Washington, D. C., 1883-)* 252, 809-17, Elrod-Erickson, M., Rould, M. A., Nekudova, L. & Pabo, C. O. (1996) *Structure (London)* 4, 1171-1180, Elrod-Erickson, M., Benson, T. E. & Pabo, C. O. (1998) *Structure (London)* 6, 451-464, Kim, C. A. & Berg, J. M. (1996) *Nature Structural Biology* 3, 940-945), though variation in helical presentation can allow for recognition of a more extended site (Pavletich, N. P. & Pabo, C. O. (1993) *Science (Washington, D. C., 1883-)* 261, 1701-7, Houbaviy, H. B., Usheva, A., Shenk, T. & Burley, S. K. (1996) *Proc Natl Acad Sci U S A* 93, 13577-82, Fairall, L., Schwabe, J. W. R., Chapman, L., Finch, J. T. & Rhodes, D. (1993) *Nature (London)* 366, 483-7, Wuttke, D. S., Foster, M. P., Case, D. A., Gottesfeld, J. M. & Wright, P. E. (1997) *J. Mol. Biol.* 273, 183-

206). In contrast to most transcription factors that rely on dimerization of protein domains for extending protein-DNA contacts to longer DNA sequences or addresses, simple covalent tandem repeats of the zinc finger domain allow for the recognition of longer asymmetric sequences of DNA by this motif.

We have recently described polydactyl zinc finger proteins that contain 6 zinc finger domains and bind 18 base pairs of contiguous DNA sequence (Liu, Q., Segal, D. J., Ghiara, J. B. & Barbas III, C. F. (1997) *PNAS* 94, 5525-5530). Recognition of 18 bps of DNA is sufficient to describe a unique DNA address within all known genomes, a requirement for using polydactyl proteins as highly specific gene switches. Indeed, control of both gene activation and repression has been shown using these polydactyl proteins in a model system (Liu, Q., Segal, D. J., Ghiara, J. B. & Barbas III, C. F. (1997) *PNAS* 94, 5525-5530).

Since each zinc finger domain typically binds three base pairs of sequence, a complete recognition alphabet requires the characterization of 64 domains. Existing information which could guide the construction of these domains has come from three types of studies:

structure determination (Pavletich, N. P. & Pabo, C. O. (1991) *Science (Washington, D. C., 1883-)* 252, 809-17, Elrod-Erickson, M., Rould, M. A., Neklodova, L. & Pabo, C. O. (1996) *Structure (London)* 4, 1171-1180, Elrod-Erickson, M., Benson, T. E. & Pabo, C. O. (1998) *Structure (London)* 6, 451-464, Kim, C. A. & Berg, J. M. (1996) *Nature Structural Biology* 3, 940-945, Pavletich, N. P. & Pabo, C. O. (1993) *Science (Washington, D. C., 1883-)* 261, 1701-7, Houbavii, H. B., Usheva, A., Shenk, T. & Burley, S. K. (1996) *Proc Natl Acad Sci U S A* 93, 13577-82, Falrall, L., Schwabe, J. W. R., Chapman, L., Finch, J. T. & Rhodes, D. (1993) *Nature (London)* 365, 483-7., 11, Wuttke, D. S., Foster, M. P., Case, D. A., Gottesfeld, J. M. & Wright, P. E. (1997) *J. Mol. Biol.* 273, 183-206., Nolte, R. T., Conlin, R. M., Harrison, S. C. & Brown, R. S. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 2938-2943, Narayan, V. A., Kriwacki, R. W. & Caradonna, J. P. (1997) *J. Biol. Chem.* 272, 7801-7809., site-directed mutagenesis (Isalan, M., Choo, Y. & Klug, A. (1997) *Proc. Natl. Acad. Sci. U. S. A.* 94, 5617-5621, Nardelli, J., Gibson, T. J., Vesque, C. & Charnay, P. (1991) *Nature* 349, 175-178, Nardelli, J., Gibson, T. & Charnay, P. (1992) *Nucleic Acids Res.* 20, 4137-44, Taylor, W. E., Suruki, H. K., Lin, A. H. T., Naraghi-Arani, P., Igarashi, R. Y., Younessian, M., Katkus, P. & Vo, N. V. (1995) *Biochemistry* 34, 3222-3230, Desjarlais, J. R. & Berg, J. M. (1992) *Proteins: Struct., Funct., Genet.* 12, 101-4, Desjarlais, J. R. & Berg, J. M. (1992) *Proc Natl Acad Sci U S A* 89, 7345-9, and phage-display selections (Choo, Y. & Klug, A. (1994) *Proc Natl Acad Sci U S A* 91, 11163-7, Greisman, H. A. & Pabo, C. O.

(1997) *Science (Washington, D. C.)* 275, 657-661.23, Rebar, E. J. & Pabo, C. O. (1994) *Science (Washington, D. C., 1883-)* 263, 671-3, Jamieson, A. C., Kim, S.-H. & Wells, J. A. (1994) *Biochemistry* 33, 5689-5695, Jamieson, A. C., Wang, H. & Kim, S.-H. (1996) *PNAS* 93, 12834-12839, Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* 37, 12026-33, Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *PNAS* 92, 344-348). All have contributed significantly to our understanding of zinc finger/DNA recognition, but each has its limitations. Structural studies have identified a diverse spectrum of protein/DNA interactions but do not explain if alternative interactions might be more optimal. Further, while interactions that allow for sequence specific recognition are observed, little information is provided on how alternate sequences are excluded from binding. These questions have been partially addressed by mutagenesis of existing proteins, but the data is always limited by the number of mutants that can be characterized. Phage-display and selection of randomized libraries overcomes certain numerical limitations, but providing the appropriate selective pressure to ensure that both specificity and affinity drive the selection is difficult. Experimental studies from several laboratories (Choo, Y. & Klug, A. (1994) *Proc Natl Acad Sci U S A* 91, 11163-7, Greisman, H. A. & Pabo, C. O. (1997) *Science (Washington, D. C.)* 275, 657-661, Rebar, E. J. & Pabo, C. O. (1994) *Science (Washington, D. C., 1883-)* 263, 671-3, Jamieson, A. C., Kim, S.-H. & Wells, J. A. (1994) *Biochemistry* 33, 5689-5695.25, Jamieson, A. C., Wang, H. & Kim, S.-H. (1996) *PNAS* 93, 12834-12839, Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* 37, 12026-33), including our own (Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *PNAS* 92, 344-348), have demonstrated that it is possible to design or select a few members of this recognition alphabet. However, the specificity and affinity of these domains for their target DNA was rarely investigated in a rigorous and systematic fashion in these early studies.

Since Jacob and Monod questioned the chemical nature of the repressor and proposed a scheme by which the synthesis of individual proteins within a cell might be provoked or repressed, specific experimental control of gene expression has been a tantalizing prospect (Jacob, F. & Monod, J. (1961) *J. Mol. Biol.* 3, 318-356). It is now well established that genomes are regulated at the level of transcription primarily through the action of proteins known as transcription factors that bind DNA in a sequence specific fashion. Often these protein factors act in a complex combinatorial manner allowing temporal, spatial, and environmentally-responsive control of gene expression (Ptashne, M. (1997) *Nature Medicine* 3, 1069-1072). Transcription factors frequently act both through a

5 DNA-binding domain which localizes the protein to a specific site within the genome, and
through accessory effector domains which act to provoke (activate) or repress transcription
10 at or near that site (Cowell, I. G. (1994) *Trends Biochem. Sci.* 19, 38-42). Effector domains,
such as the activation domain VP16 (Sadowski, I., Ma, J., Triezenberg, S. & Ptashne, M.
(1988) *Nature* 335, 563-564) and the repression domain KRAB (Margolin, J. F., Friedman,
15 J. R., Meyer, W., K.-H., Vissing, H., Thiesen, H.-J. & Rauscher III, F. J. (1994) *Proc. Natl.
Acad. Sci. USA* 91, 4509-4513), are typically modular and retain their activity when they are
fused to other DNA-binding proteins. Whereas genes might be readily controlled by
directing transcription factors to particular sites within a genome, the design of DNA binding
20 proteins that might be fashioned to bind any given sequence has been a daunting
challenge.

The present disclosure is based on the recognition of the structural features unique to
the Cys₂-His₂ class of nucleic acid-binding, zinc finger proteins. The Cys₂-His₂ zinc finger
25 domain consists of a simple $\beta\alpha$ fold of approximately 30 amino acids in length. Structural
stability of this fold is achieved by hydrophobic interactions and by chelation of a single zinc
ion by the conserved Cys₂-His₂ residues (Lee, M. S., Gippert, G. P., Soman, K. V., Case, D.
A. & Wright, P. E. (1989) *Science* 245, 635-637). Nucleic acid recognition is achieved
30 through specific amino acid side chain contacts originating from the α -helix of the domain,
which typically binds three base pairs of DNA sequence (Pavletich, N. P. & Pabo, C. O.
(1991) *Science* 252, 809-17, Elrod-Erickson, M., Rould, M. A., Neklodova, L. & Pabo, C. O.
(1996) *Structure* 4, 1171-1180). Unlike other nucleic acid recognition motifs, simple covalent
35 linkage of multiple zinc finger domains allows the recognition of extended asymmetric
sequences of DNA. Studies of natural zinc finger proteins have shown that three zinc finger
domains can bind 9 bp of contiguous DNA sequence (Pavletich, N. P. & Pabo, C. O. (1991)
40 *Science* 252, 809-17., Swirnoff, A. H. & Milbrandt, J. (1995) *Mol. Cell. Biol.* 15, 2275-87).
Whereas recognition of 9 bp of sequence is insufficient to specify a unique site within even
the small genome of *E. coli*, polydactyl proteins containing six zinc finger domains can
specify 18-bp recognition (Liu, Q., Segal, D. J., Ghilardi, J. B. & Barbas III, C. F. (1997) *Proc.*
45 *Natl. Acad. Sci. USA* 94, 5525-5530). With respect to the development of a universal
system for gene control, an 18-bp address can be sufficient to specify a single site within all
known genomes. While polydactyl proteins of this type are unknown in nature, however,
50 their efficacy in gene activation and repression within living human cells has recently been

shown (Liu, Q., Segal, D. J., Ghiara, J. B. & Barbas III, C. F. (1997) *Proc. Natl. Acad. Sci. USA* 94, 5525-5530).

Brief Summary of the Invention

In one aspect, the present invention provides an isolated and purified zinc finger-nucleotide binding polypeptide that contains the amino acid residue sequence of any of SEQ ID NO:1-16. In a related aspect, this invention further provides compositions comprising from two to about 12 such zinc finger-nucleotide binding polypeptides. The composition preferably contains from 2 to about 6 polypeptides. In a preferred embodiment, the zinc finger-nucleotide binding polypeptide are operatively linked, preferably by an amino acid residue linker having the sequence of SEQ ID NO 111. A composition of this invention specifically binds a nucleotide target that contains the sequence 5'-(GNN)_n-3', wherein each N is A, C, G, or T with the proviso that all N's cannot be C and where n is preferably 2 to 6. A polypeptide or composition can be further operatively linked to one or more transcription modulating factors such as a transcription activators or transcription suppressors or repressors. The present invention also provides an isolated and purified polynucleotide that encodes a polypeptide or composition of this invention and an expression vector containing such a polynucleotide.

In a still further aspect, the present invention provides a process of regulating the function of a nucleotide sequence that contains the sequence 5'-(GNN)_n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of a composition of this invention operatively linked to one or more transcription modulating factors. The 5'-(GNN)_n-3' sequence can be found in the transcribed region or promotor region of the nucleotide or within an expressed sequence tag.

The present disclosure demonstrates the simplicity and efficacy of a general strategy for the rapid production of gene switches. With a family of defined zinc finger domains recognizing sequences of the 5'-GNN-3' subset of a 64 member zinc finger alphabet, polydactyl proteins specifically recognizing novel 9- or, for the first time, 18-bp sequences were constructed and characterized. Potent transcription factors were generated and shown to control both gene activation and repression. Gene activation was achieved using the herpes simplex virus VP16 activation domain (Sadowski, I., Ma, J., Triezenberg, S. & Ptashne, M. (1988) *Nature* 335, 563-564) and a recombinant tetrameric repeat of its minimal activation domain. Gene repression or silencing was achieved using three effector

5 domains of human origin, the *krüppel* associated box (KRAB) (Margolin, J. F., Friedman, J. R., Meyer, W., K.-H., Vissing, H., Thiesen, H.-J. & Rauscher III, F. J. (1994) *Proc. Natl.*
10 *Acad. Sci. USA* **91**, 4509-4513), the ERF repressor domain (ERD) (Sgouras, D. N., Athanasiou, M. A., Beal, G. J., Jr., Fisher, R. J., Blair, D. G. & Mavrothalassitis, G. J. (1995) *EMBO J.* **14**, 4781-4793), and the mSIN3 interaction domain (SID) (Ayer, D. E., Laherty, C. D., Lawrence, Q. A., Armstrong, A. P. & Eisenman, R. N. (1996) *Mol. Cell. Biol.* **16**, 5772-
15 5781). Using luciferase reporter gene assays in human epithelial cells, the data show that artificial transcriptional regulators, designed to target the promoter of the proto-oncogene *erbB-2/HER-2*, can ablate or activate gene expression in a specific manner. For the first
20 time, gene activation or repression was achieved by targeting within the gene transcript, suggesting that information obtained from expressed sequence tags (ESTs) may be sufficient for the construction of gene switches. The novel methodology and materials described herein promise diverse applications in gene therapy, transgenic organisms, functional genomics, and other areas of cell and molecular biology.

Brief Description of the Drawing

In the drawing, which forms a portion of the specification

30 FIG. 1 (shown in six panels) shows the binding specificity of regions of zinc finger-nucleotide binding polypeptides of the invention.

Detailed Description of the Invention

I. The Invention

35 The present invention provides zinc finger-nucleotide binding polypeptides, compositions containing one or more such polypeptides and the use of the polypeptides and compositions for modulating gene expression.

II Compounds

40 A compound of this invention is an isolated zinc finger-nucleotide binding polypeptide that binds to a GNN nucleotide sequence and modulates the function of that nucleotide sequence. The polypeptide can enhance or suppress transcription of a gene, and can bind to DNA or RNA. A zinc finger-nucleotide binding polypeptide refers to a polypeptide which is a derivatized form of a wild-type zinc finger protein or one produced
45 through recombination. A polypeptide may be a hybrid which contains zinc finger domain(s)
50
55

5 from one protein linked to zinc finger domain(s) of a second protein, for example. The
domains may be wild type or mutagenized. A polypeptide includes a truncated form of a
10 wild type zinc finger protein. Examples of zinc finger proteins from which a polypeptide can
be produced include TFIIIA and zif268.

A zinc finger-nucleotide binding polypeptide of this invention comprises a unique
15 heptamer (contiguous sequence of 7 amino acid residues) within the α -helical domain of the
polypeptide, which heptameric sequence determines binding specificity to a target
nucleotide. That heptameric sequence can be located anywhere within the α -helical
domain but it is preferred that the heptamer extend from position -1 to position 6 as the
20 residues are conventionally numbered in the art. A polypeptide of this invention can include
any β -sheet and framework sequences known in the art to function as part of a zinc finger
protein. A large number of zinc finger-nucleotide binding polypeptides were made and
tested for binding specificity against target nucleotides containing a GNN triplet. The results
25 of those studies are summarized in FIG. 1. In FIG. 1, the GNN triplet binding specificity for
each peptide is shown in the right-hand column, with the highest specificity shown first and
in boldface. In FIG. 1, SEQ ID Nos. are shown in parentheses. For each particular GNN
(e.g., GAA, shown in the right-hand column of FIG. 1) target, the sequences are listed in
30 order of decreasing specificity for that Triplet.

As shown in FIG. 1, the data show a striking conservation of all three of the primary
DNA contact positions (-1, 3, and 6) was observed for virtually all the clones of a given
35 target. Although many of these residues were observed previously at these positions
following selections with much less complete libraries, the extent of conservation observed
here represents a dramatic improvement over earlier studies (Choo, Y. & Klug, A. (1994)
Proc Natl Acad Sci U S A 91, 11163-7, Greisman, H. A. & Pabo, C. O. (1997) *Science*
(Washington, D. C.) 275, 657-661, Rebar, E. J. & Pabo, C. O. (1994) *Science*
40 (Washington, D. C., 1883-) 263, 671-3, Jamieson, A. C., Kim, S.-H. & Wells, J. A. (1994)
Biochemistry 33, 5689-5695, Jamieson, A. C., Wang, H. & Kim, S.-H. (1996) *PNAS* 93,
12834-12839, Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *PNAS* 92, 344-348). The
45 present invention discloses that the teachings of the prior art that the three helical positions
-1, 3, and 6 of a zinc finger domain are sufficient to allow for the detailed description of the
DNA binding specificity of the domain are incorrect.

Typically, phage selections have shown a consensus selection in only one or two of
50 these positions. The greatest sequence variation occurred at the residues in positions 1

5 and 5, which do not make bases contacts in the Zif268/DNA structure and were expected
not to contribute significantly to recognition (Pavletich, N. P. & Pabo, C. O. (1991) *Science*
10 (*Washington, D. C., 1883-*) 252, 809-17, Elrod-Erickson, M., Rould, M. A., Nekludova, L. &
Pabo, C. O. (1996) *Structure (London)* 4, 1171-1180). Variation in positions 1 and 5 also
implied that the conservation in the other positions was due to their interaction with the DNA
and not simply the fortuitous amplification of a single clone due to other reasons.

15 Conservation of residue identity at position 2 was also observed. The conservation of
position -2 is somewhat artifactual; the NNK library had this residue fixed as serine. This
residue makes contacts with the DNA backbone in the Zif268 structure. Both libraries
contained an invariant leucine at position 4, a critical residue in the hydrophobic core that
20 stabilizes folding of this domain.

Impressive amino acid conservation was observed for recognition of the same
nucleotide in different targets. For example, Asn in position 3 (Asn3) was virtually always
25 selected to recognize adenine in the middle position, whether in the context of GAG, GAA,
GAT, or GAC. Gln-1 and Arg-1 were always selected to recognize adenine or guanine,
respectively, in the 3' position regardless of context. Amide side chain based recognition of
adenine by Gln or Asn is well documented in structural studies as is the Arg guanidinium
30 side chain to guanine contact with a 3' or 5' guanine (Elrod-Erickson, M., Benson, T. E. &
Pabo, C. O. (1998) *Structure (London)* 6, 451-464, Kim, C. A. & Berg, J. M. (1996) *Nature*
Structural Biology 3, 940-945., Fairall, L., Schwabe, J. W. R., Chapman, L., Finch, J. T. &
Rhodes, D. (1993) *Nature (London)* 366, 483-7). More often, however, two or three amino
35 acids were selected for nucleotide recognition. His3 or Lys3 (and to a lesser extent, Gly3)
were selected for the recognition of a middle guanine. Ser3 and Ala3 were selected to
recognize a middle thymine. Thr3, Asp3, and Glu3 were selected to recognize a middle
cytosine. Asp and Glu were also selected in position -1 to recognize a 3' cytosine, while
40 Thr-1 and Ser-1 were selected to recognize a 3' thymine.

Selected Zif268 variants were subcloned into a bacterial expression vector, and the
proteins overexpressed (finger-2 proteins, hereafter referred to by the subsite for which they
45 were panned). It is important to study soluble proteins rather than phage-fusions since it is
known that the two may differ significantly in their binding characteristics (Cramer, A.,
Cwirla, S. & Stemmer, W. P. (1996) *Nat. Med.* 2, 100-102). The proteins were tested for
their ability to recognize each of the 16 5'-GNN-3' finger -2 subsites using a multi-target
50 ELISA assay. This assay provided an extremely rigorous test for specificity since there

5
10
15
20
25
30
35
40
45
50
55

were always six "non-specific" sites which differed from the "specific" site by only a single nucleotide out of a nine-nucleotide target. Many of the phage-selected finger-2 proteins showed exquisite specificity, while others demonstrated varying degrees of crossreactivity. Some polypeptides actually bound better to subsites other than those for which they were selected.

Attempts were made to improve binding specificity by modifying the recognition helix using site-directed mutagenesis. Data from our selections and structural information guided mutant design. As the most exhaustive study performed to date, over 100 mutant proteins were characterized in an effort to expand our understanding of the rules of recognition. Although helix positions 1 and 5 are not expected to play a direct role in DNA recognition, the best improvements in specificity always involved modifications in these positions. These residues have been observed to make phosphate backbone contacts, which contribute to affinity in a non-sequence specific manner. Removal of non-specific contacts increases the importance of the specific contacts to the overall stability of the complex, thereby enhancing specificity. For example, the specificity of polypeptides for target triplets GAC, GAA, and GAG were improved simply by replacing atypical, charged residues in positions 1 and 5 with smaller, uncharged residues.

Another class of modifications involved changes to both binding and non-binding residues. The crossreactivity of polypeptides for GGG and the finger-2 subsite GAG was abolished by the modifications His3Lys and Thr5Val. It is interesting to note that His3 was unanimously selected during panning to recognize the middle guanine, although Lys3 provided better discrimination of A and G. This suggests that panning conditions for this protein may have favored selection by a parameter such as affinity over that of specificity. In the Zif268 structure, His3 donates a hydrogen bond to the N7 of the middle guanine (Pavletich, N. P. & Pabo, C. O. (1991) *Science (Washington, D. C., 1983-)* **252**, 809-17, Elrod-Erickson, M., Rould, M. A., Nekudova, L. & Pabo, C. O. (1996) *Structure (London)* **4**, 1171-1180). This bond could also be made with N7 of adenine, and in fact Zif268 does not discriminate between G and A in this position (Swimoff, A. H. & Milbrandt, J. (1995) *Mol. Cell. Biol.* **15**, 2275-87). His3 was found to specify only a middle guanine in polypeptides targeted to GGA, GGC, and GGT, even though Lys3 was selected during panning for GGC and GGT. Similarly, the multiple crossreactivities of polypeptides targeted to GTG were attenuated by modifications Lys1Ser and Ser3Glu, resulting in a 5-fold loss in affinity. Glu3 has been shown to be very specific for cytosine in binding site selection studies of Zif268

(Swimoff, A. H. & Milbrandt, J. (1995) *Mol. Cell. Biol.* 15, 2275-87). No structural studies show an interaction of Glu3 with the middle thymine, and Glu3 was never selected to recognize a middle thymine in our study or any others (Choo, Y. & Klug, A. (1994) *Proc Natl Acad Sci U S A* 91, 11163-7, Greisman, H. A. & Pabo, C. O. (1997) *Science (Washington, D. C.)* 275, 657-661, Rebar, E. J. & Pabo, C. O. (1994) *Science (Washington, D. C., 1883-)* 263, 671-3, Jamieson, A. C., Kim, S.-H. & Wells, J. A. (1994) *Biochemistry* 33, 5689-5695, Jamieson, A. C., Wang, H. & Kim, S.-H. (1996) *PNAS* 93, 12834-12839, Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* 37, 12026-33, Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *PNAS* 92, 344-348). Despite this, the Ser3Glu modification favored the recognition of a middle thymine over cytosine. These examples illustrate the limitations of relying on previous structures and selection data to understand the structural elements underlying specificity. It should also be emphasized that improvements by modifications involving positions 1 and 5 could not have been predicted by existing "recognition codes" (Desjarlais, J. R. & Berg, J. M. (1992) *Proc Natl Acad Sci U S A* 89, 7345-9, Suzuki, M., Gerstein, M. & Yagi, N. (1994) *Nucleic Acids Res.* 22, 3397-405, Choo, Y. & Klug, A. (1994) *Proc. Natl. Acad. Sci. U. S. A.* 91, 11168-72, Choo, Y. & Klug, A. (1997) *Curr. Opin. Struct. Biol.* 7, 117-125), which typically only consider positions -1, 2, 3, and 6. Only by the combination of selection and site-directed mutagenesis can we begin to fully understand the intricacies of zinc finger/DNA recognition.

From the combined selection and mutagenesis data it emerged that specific recognition of many nucleotides could be best accomplished using motifs, rather than a single amino acid. For example, the best specification of a 3' guanine was achieved using the combination of Arg-1, Ser1, and Asp2 (the RSD motif). By using Val5 and Arg6 to specify a 5' guanine, recognition of subsites GGG, GAG, GTG, and GCG could be accomplished using a common helix structure (SRSD-X-LVR) differing only in the position 3 residue (Lys3 for GGG, Asn3 for GAG, Glu3 for GTG, and Asp3 for GCG). Similarly, 3' thymine was specified using Thr-1, Ser1, and Gly2 in the final clones (the TSG motif). Further, a 3' cytosine could be specified using Asp-1, Pro1, and Gly2 (the DPG motif) except when the subsite was GCC; Pro1 was not tolerated by this subsite. Specification of a 3' adenine was with Gln-1, Ser1, Ser2 in two clones (QSS motif). Residues of positions 1 and 2 of the motifs were studied for each of the 3' bases and found to provide optimal specificity for a given 3' base as described here.

5

10

15

20

25

30

35

40

45

50

55

The multi-target ELISA assay assumed that all the proteins preferred guanine in the 5' position since all proteins contained Arg6 and this residue is known from structural studies to contact guanine at this position (Pavletich, N. P. & Pabo, C. O. (1991) *Science* (Washington, D. C., 1983-) 252, 809-17, Elrod-Erickson, M., Rould, M. A., Nekludova, L. & Pabo, C. O. (1996) *Structure* (London) 4, 1171-1180, Elrod-Erickson, M., Benson, T. E. & Pabo, C. O. (1998) *Structure* (London) 6, 451-464, Kim, C. A. & Berg, J. M. (1996) *Nature Structural Biology* 3, 940-945, Pavletich, N. P. & Pabo, C. O. (1993) *Science* (Washington, D. C., 1983-) 261, 1701-7, Houbaviy, H. B., Usheva, A., Shenk, T. & Burley, S. K. (1996) *Proc Natl Acad Sci U S A* 93, 13577-82, Fairall, L., Schwabe, J. W. R., Chapman, L., Finch, J. T. & Rhodes, D. (1993) *Nature* (London) 366, 483-7, Wuttke, D. S., Foster, M. P., Case, D. A., Gottesfeld, J. M. & Wright, P. E. (1997) *J. Mol. Biol.* 273, 183-206, Nolte, R. T., Conlin, R. M., Harrison, S. C. & Brown, R. S. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 2938-2943). This interaction was demonstrated here using the 5' binding site signature assay ((Choo, Y. & Klug, A. (1994) *Proc. Natl. Acad. Sci. U. S. A.* 91, 11168-72); Fig. 2, white bars). Each protein was applied to pools of 16 oligonucleotide targets in which the 5' nucleotide of the finger-2 subsite was fixed as G, A, T, or C and the middle and 3' nucleotides were randomized. All proteins preferred the GNN pool with essentially no crossreactivity.

The results of the multi-target ELISA assay were confirmed by affinity studies of purified proteins. In cases where crossreactivity was minimal in the ELISA assay, a single nucleotide mismatch typically resulted in a greater than 100-fold loss in affinity. This degree of specificity had yet to be demonstrated with zinc finger proteins. In general, proteins selected or designed to bind subsites with G or A in the middle and 3' position had the highest affinity, followed by those which had only one G or A in the middle or 3' position, followed by those which contained only T or C. The former group typically bound their targets with a higher affinity than Zif268 (10 nM), the latter with somewhat lower affinity, and almost all the proteins had an affinity lower than that of the parental C7 protein. There was no correlation between binding affinity and binding specificity suggesting that specificity can result not only from specific protein-DNA contacts, but also from interactions which exclude all but the correct nucleotide.

Asp2 was always co-selected with Arg-1 in all proteins for which the target subsite was GNG. It is now understood that there are two reasons for this. From structural studies of Zif268 (Pavletich, N. P. & Pabo, C. O. (1991) *Science* (Washington, D. C., 1983-) 252,

809-17, Elrod-Erickson, M., Rould, M. A., Nekludova, L. & Pabo, C. O. (1996) *Structure (London)* **4**, 1171-1180), it is known that Asp2 of finger 2 makes a pair of buttressing hydrogen bonds with Arg-1 which stabilize the Arg-1/3' guanine interaction, as well as some water-mediated contacts. However, the carboxylate of Asp2 also accepts a hydrogen bond from the N4 of a cytosine that is base-paired to a 5' guanine of the finger-1 subsite. Adenine base paired to T in this position can make an analogous contact to that seen with cytosine. This interaction is particularly important because it extends the recognition subsite of finger 2 from three nucleotides (GNG) to four (GNG(G/T)) (Isalan, M., Choo, Y. & Klug, A. (1997) *Proc. Natl. Acad. Sci. U. S. A.* **94**, 5617-5621., Jamieson, A. C., Wang, H. & Kim, S.-H. (1996) *PNAS* **93**, 12834-12839, Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* **37**, 12026-33). This phenomenon is referred to as "target site overlap", and has three important ramifications. First, Asp2 was favored for selection by our library when the finger-2 subsite was GNG because our finger-1 subsite contained a 5' guanine. Second, it may limit the utility of the libraries used in this study to selection on GNN or TNN finger-2 subsites because finger 3 of these libraries contains an Asp2, which may help specify the 5' nucleotide of the finger-2 subsite to be G or T. In Zif268 and C7, which have Thr6 in finger 2, Asp2 of finger 3 enforces G or T recognition in the 5' position (T/G)GG. This interaction may also explain why previous phage display studies, which all used Zif268-based libraries, have found selection limited primarily to GNN recognition (Choo, Y. & Klug, A. (1994) *Proc Natl Acad Sci U S A* **91**, 11163-7., Rebar, E. J. & Pabo, C. O. (1994) *Science (Washington, D. C., 1883-)* **263**, 671-3, Jamieson, A. C., Kim, S.-H. & Wells, J. A. (1994) *Biochemistry* **33**, 5689-5695, Jamieson, A. C., Wang, H. & Kim, S.-H. (1996) *PNAS* **93**, 12834-12839, Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* **37**, 12026-33, Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *PNAS* **92**, 344-348).

Finally, target site overlap potentially limits the use of these zinc fingers as modular building blocks. From structural data it is known that there are some zinc fingers in which target site overlap is quite extensive, such as those in GLI and YY1, and others which are similar to Zif268 and display only modest overlap. In our final set of proteins, Asp2 is found in polypeptides that bind GGG, GAG, GTG, and GCG. The overlap potential of other residues found at position 2 is largely unknown, however structural studies reveal that many other residues found at this position may participate in such cross-subsite contacts. Fingers containing Asp2 may limit modularity, since they would require that each GNG subsite be followed by a T or G.

Table 1, below, summarized the sequences (SEQ ID NOs;1-16) showing the highest selectivity for the sixteen embodiment of GNN target triplets.

Table 1

Target Specificity	amino acids positions -1 1 2 3 4 5 6	SEQ ID NO:
GAA	QSSNLVR	1
GAC	DPGNLVR	2
GAG	RSDNLVR	3
GAT	TSGNLVR	4
GCA	QSGDLRR	5
GCC	DCRDLAR	6
GCG	RSDDLVR	7
GCT	TSGELVR	8
GGA	QRAHLER	9
GGC	DPGHLVR	10
GGG	RSDKLVR	11
GGT	TSGHLVR	12
GTA	QSSSLVR	13
GTC	DPGALVR	14
GTG	RSDELVR	15
GTT	TSGSLVR	16

The data show that all possible GNN triplet sequences can be recognized with exquisite specificity by zinc finger domains. Optimized zinc finger domains can discriminate single base differences by greater than 100-fold loss in affinity. While many of the amino acids found in the optimized proteins at the key contact positions -1,3, and 6 are those that are consistent with a simple code of recognition, it has been discovered that

5 optimal specific recognition is sensitive to the context in which these residues are presented. Residues at positions 1,2, and 5 have been found to be critical for specific
10 recognition. Further the data demonstrates for the first time that sequence motifs at positions -1,1, and 2 rather than the simple identity of the position 1 residue are required for highly specific recognition of the 3' base. These residues likely provide the proper stereochemical context for interactions of the helix both in terms of recognition of specific
15 bases and in the exclusion of other bases, the net result being highly specific interactions. Broad utility of these domains would be realized if they were modular in both their interactions with DNA and other zinc finger domains. This could be achieved by working within the likely limitations imposed by target site overlap, namely that sequences of the 5'-
20 (GNN)_n-3' type should be targeted. Ready recombination of the disclosed domains then allows for the creation of polydactyl proteins of defined specificity precluding the need to develop phage display libraries in their generation. These polydactyl proteins have been used to activate and repress transcription driven by the human *erbB-2* promoter in living
25 cells. The family of zinc finger domains described herein is likely sufficient for the construction of 16⁶ or 17 million novel proteins that bind the 5'-(GNN)_n-3' family of DNA sequences.

30 The zinc finger-nucleotide binding polypeptide derivative can be derived or produced from a wild type zinc finger protein by truncation or expansion, or as a variant of the wild type-derived polypeptide by a process of site directed mutagenesis, or by a combination of the procedures. The term "truncated" refers to a zinc finger-nucleotide binding polypeptide that contains less than the full number of zinc fingers found in the native zinc finger binding
35 protein or that has been deleted of non-desired sequences. For example, truncation of the zinc finger-nucleotide binding protein TFIIIA, which naturally contains nine zinc fingers, might be a polypeptide with only zinc fingers one through three. Expansion refers to a zinc finger polypeptide to which additional zinc finger modules have been added. For example,
40 TFIIIA may be extended to 12 fingers by adding 3 zinc finger domains. In addition, a truncated zinc finger-nucleotide binding polypeptide may include zinc finger modules from more than one wild type polypeptide, thus resulting in a "hybrid" zinc finger-nucleotide
45 binding polypeptide.

The term "mutagenized" refers to a zinc finger derived-nucleotide binding polypeptide that has been obtained by performing any of the known methods for accomplishing random or site-directed mutagenesis of the DNA encoding the protein. For
50 instance, in TFIIIA, mutagenesis can be performed to replace nonconserved residues in one

5

or more of the repeats of the consensus sequence. Truncated zinc finger-nucleotide binding proteins can also be mutagenized.

10

Examples of known zinc finger-nucleotide binding polypeptides that can be truncated, expanded, and/or mutagenized according to the present invention in order to inhibit the function of a nucleotide sequence containing a zinc finger-nucleotide binding motif includes TFIIIA and zif268. Other zinc finger-nucleotide binding proteins will be known to those of skill in the art.

15

20

A polypeptide of this invention can be made using a variety of standard techniques well known in the art (See, e.g., United States Patent Application No. 08/676,318, filed 1/18/1995, the entire disclosure of which is incorporated herein by reference). Phage display libraries of zinc finger proteins were created and selected under conditions that favored enrichment of sequence specific proteins. Zinc finger domains recognizing a number of sequences required refinement by site-directed mutagenesis that was guided by both phage selection data and structural information.

25

30

35

40

45

50

55

The murine Cys₂-His₂ zinc finger protein Zif268 is used for construction of phage display libraries (Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *PNAS* 92, 344-348). Zif268 is structurally the most well characterized of the zinc-finger proteins (Pavletich, N. P. & Pabo, C. O. (1991) *Science (Washington, D. C., 1883-)* 252, 809-17, Elrod-Erickson, M., Rould, M. A., Neklodova, L. & Pabo, C. O. (1996) *Structure (London)* 4, 1171-1180, Swirnoff, A. H. & Milbrandt, J. (1995) *Mol. Cell. Biol.* 15, 2275-87). DNA recognition in each of the three zinc finger domains of this protein is mediated by residues in the N-terminus of the α -helix contacting primarily three nucleotides on a single strand of the DNA. The operator binding site for this three finger protein is 5'-GCGTGGGCG-3' (finger-2 subsite is underlined). Structural studies of Zif268 and other related zinc finger-DNA complexes (Elrod-Erickson, M., Benson, T. E. & Pabo, C. O. (1998) *Structure (London)* 6, 451-464, Kim, C. A. & Berg, J. M. (1996) *Nature Structural Biology* 3, 940-945, Pavletich, N. P. & Pabo, C. O. (1993) *Science (Washington, D. C., 1883-)* 261, 1701-7, Houbaviy, H. B., Usheva, A., Shenk, T. & Burley, S. K. (1996) *Proc Natl Acad Sci U S A* 93, 13577-82, Fairall, L., Schwabe, J. W. R., Chapman, L., Finch, J. T. & Rhodes, D. (1993) *Nature (London)* 366, 483-7, Wuttke, D. S., Foster, M. P., Case, D. A., Gottesfeld, J. M. & Wright, P. E. (1997) *J. Mol. Biol.* 273, 183-206., Nolte, R. T., Conlin, R. M., Harrison, S. C. & Brown, R. S. (1998) *Proc. Natl. Acad. Sci. U. S. A.* 95, 2938-2943, Narayan, V. A., Kriwacki, R. W. & Caradonna, J. P. (1997) *J. Biol. Chem.* 272, 7801-7809) have shown that residues from

5 primarily three positions on the α -helix, -1, 3, and 6, are involved in specific base contacts. Typically, the residue at position -1 of the α -helix contacts the 3' base of that finger's subsite
10 while positions 3 and 6 contact the middle base and the 5' base, respectively.

In order to select a family of zinc finger domains recognizing the 5'-GNN-3' subset of sequences, two highly diverse zinc finger libraries were constructed in the phage display vector pComb3H (Barbas III, C. F., Kang, A. S., Lerner, R. A. & Benkovic, S. J. (1991) *Proc. Natl. Acad. Sci. USA* 88, 7978-7982., Rader, C. & Barbas III, C. F. (1997) *Curr. Opin. Biotechnol.* 8, 503-508). Both libraries involved randomization of residues within the α -helix of finger 2 of C7, a variant of Zif268 (Wu, H., Yang, W.-P. & Barbas III, C. F. (1995) *PNAS* 92, 344-348). Library 1 was constructed by randomization of positions -1,1,2,3,5,6 using a NNK doping strategy while library 2 was constructed using a VNS doping strategy with randomization of positions -2,-1,1,2,3,5,6. The NNK doping strategy allows for all amino acid combinations within 32 codons while VNS precludes Tyr, Phe, Cys and all stop codons in its 24 codon set. The libraries consisted of 4.4×10^9 and 3.5×10^9 members, respectively,
20 each capable of recognizing sequences of the 5'-GCGNNGCG-3' type. The size of the NNK library ensured that it could be surveyed with 99% confidence while the VNS library was highly diverse but somewhat incomplete. These libraries are, however, significantly larger than previously reported zinc finger libraries (Choo, Y. & Klug, A. (1994) *Proc Natl Acad Sci U S A* 91, 11163-7, Greisman, H. A. & Pabo, C. O. (1997) *Science (Washington, D. C.)* 275, 657-661, Rebar, E. J. & Pabo, C. O. (1994) *Science (Washington, D. C., 1883-)* 263, 671-3, Jamieson, A. C., Kim, S.-H. & Wells, J. A. (1994) *Biochemistry* 33, 5689-5695,
30 Jamieson, A. C., Wang, H. & Kim, S.-H. (1996) *PNAS* 93, 12834-12839, Isalan, M., Klug, A. & Choo, Y. (1998) *Biochemistry* 37, 12028-33). Seven rounds of selection were performed on the zinc finger displaying-phage with each of the 16 5'-GCGNNGCG-3' biotinylated hairpin DNAs targets using a solution binding protocol. Stringency was increased in each round by the addition of competitor DNA. Sheared herring sperm DNA was provided for selection against phage that bound non-specifically to DNA. Stringent selective pressure for sequence specificity was obtained by providing DNAs of the 5'-GCGNNGCG-3' types as specific competitors. Excess DNA of the 5'-GCGNNGCG-3' type was added to provide even more stringent selection against binding to DNAs with single or double base changes as compared to the biotinylated target. Phage binding to the single biotinylated DNA target sequence were recovered using streptavidin coated beads.
40 In some cases the selection process was repeated. The present data show that these

5 domains are functionally modular and can be recombined with one another to create
polydactyl proteins capable of binding 18-bp sequences with subnanomolar affinity. The
10 family of zinc finger domains described herein is sufficient for the construction of 17 million
novel proteins that bind the 5'-(GNN)₆-3' family of DNA sequences.

The invention includes a nucleotide sequence encoding a zinc finger-nucleotide
binding polypeptide. DNA sequences encoding the zinc finger-nucleotide binding
15 polypeptides of the invention, including native, truncated, and expanded polypeptides, can
be obtained by several methods. For example, the DNA can be isolated using hybridization
procedures which are well known in the art. These include, but are not limited to: (1)
20 hybridization of probes to genomic or cDNA libraries to detect shared nucleotide
sequences; (2) antibody screening of expression libraries to detect shared structural
features; and (3) synthesis by the polymerase chain reaction (PCR). RNA sequences of the
invention can be obtained by methods known in the art (See for example, Current Protocols
25 in Molecular Biology Ausubel, et al. Eds., 1989).

The development of specific DNA sequences encoding zinc finger-nucleotide
binding polypeptides of the invention can be obtained by: (1) isolation of a double-stranded
DNA sequence from the genomic DNA; (2) chemical manufacture of a DNA sequence to
30 provide the necessary codons for the polypeptide of interest; and (3) in vitro synthesis of a
double-stranded DNA sequence by reverse transcription of mRNA isolated from a
eukaryotic donor cell. In the latter case, a double-stranded DNA complement of mRNA is
eventually formed which is generally referred to as cDNA. Of these three methods for
35 developing specific DNA sequences for use in recombinant procedures, the isolation of
genomic DNA is the least common. This is especially true when it is desirable to obtain the
microbial expression of mammalian polypeptides due to the presence of introns.

For obtaining zinc finger derived-DNA binding polypeptides, the synthesis of DNA
40 sequences is frequently the method of choice when the entire sequence of amino acid
residues of the desired polypeptide product is known. When the entire sequence of amino
acid residues of the desired polypeptide is not known, the direct synthesis of DNA
sequences is not possible and the method of choice is the formation of cDNA sequences.
45 Among the standard procedures for isolating cDNA sequences of interest is the formation of
plasmid-carrying cDNA libraries which are derived from reverse transcription of mRNA
which is abundant in donor cells that have a high level of genetic expression. When used in
combination with polymerase chain reaction technology, even rare expression products can
50 be clones. In those cases where significant portions of the amino acid sequence of the

5

10

polypeptide are known, the production of labeled single or double-stranded DNA or RNA probe sequences duplicating a sequence putatively present in the target cDNA may be employed in DNA/DNA hybridization procedures which are carried out on cloned copies of the cDNA which have been denatured into a single-stranded form (Jay, et al., *Nucleic Acid Research* 11:2325, 1983).

15

In another aspect, the present invention provides a pharmaceutical composition comprising a therapeutically effective amount of a zinc finger-nucleotide binding polypeptide or a therapeutically effective amount of a nucleotide sequence that encodes a zinc finger-nucleotide binding polypeptide in combination with a pharmaceutically acceptable carrier.

20

As used herein, the terms "pharmaceutically acceptable", "physiologically tolerable" and grammatical variations thereof, as they refer to compositions, carriers, diluents and reagents, are used interchangeable and represent that the materials are capable of administration to or upon a human without the production of undesirable physiological effects such as nausea, dizziness, gastric upset and the like which would be to a degree that would prohibit administration of the composition.

25

30

The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art. Typically such compositions are prepared as sterile injectables either as liquid solutions or suspensions, aqueous or non-aqueous, however, solid forms suitable for solution, or suspensions, in liquid prior to use can also be prepared. The preparation can also be emulsified.

35

40

The active ingredient can be mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient and in amounts suitable for use in the therapeutic methods described herein. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, as well as pH buffering agents and the like which enhance the effectiveness of the active ingredient.

45

50

The therapeutic pharmaceutical composition of the present invention can include pharmaceutically acceptable salts of the components therein. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, mandelic and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium,

55

5 potassium, ammonium, calcium or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

10 Physiologically tolerable carriers are well known in the art. Exemplary of liquid carriers are sterile aqueous solutions that contain no materials in addition to the active ingredients and water, or contain a buffer such as sodium phosphate at physiological pH value, physiological saline or both, such as phosphate-buffered saline. Still further,
15 aqueous carriers can contain more than one buffer salt, as well as salts such as sodium and potassium chlorides, dextrose, propylene glycol, polyethylene glycol and other solutes. Liquid compositions can also contain liquid phases in addition to and to the exclusion of water. Exemplary of such additional liquid phases are glycerin, vegetable oils such as
20 cottonseed oil, organic esters such as ethyl oleate, and water-oil emulsions.

III. Compositions

25 In another aspect, the present invention provides a plurality of zinc finger-nucleotide binding polypeptides operatively linked in such a manner to specifically bind a nucleotide target motif defined as 5'-(GNN)*n*-3', where *n* is an Integer greater than 1. Preferably, *n* is an integer from 2 to about 6.

30 Means for linking zinc finger-nucleotide binding polypeptide are described hereinafter in the Examples as well as in United States Patent Application No. 08/676,318, filed 1/18/1995). The individual polypeptides are preferably linked with oligopeptide linkers. Such linkers preferably resemble the linker that are found in naturally occurring zinc finger
35 proteins. A preferred linker for use in the present invention is the amino acid residue sequence TGEKP (SEQ ID NO:111).

To examine the efficacy of making such compositions and their use in gene control,
40 the human *erbB-2* gene was chosen as a model. A polydactyl protein specifically recognizing an 18bp sequence in the 5'-untranslated region of this gene was converted into a transcriptional repressor by fusion with KRAB, ERD, or SID repressor domains. Transcriptional activators were generated by fusion with the herpes simplex VP16 activation domain or with a tetrameric repeat of VP16's minimal activation domain, termed VP64. The
45 data show for the first time that both gene repression and activation can be achieved by targeting designed proteins to a single site within the transcribed region of a gene.

The human *erbB-2* gene was chosen as a model target for the development of zinc
50 finger-based transcriptional switches. Members of the ErbB receptor family play important roles in the development of human malignancies. In particular, *erbB-2* is overexpressed as

5 a result of gene amplification and/or transcriptional deregulation in a high percentage of
human adenocarcinomas arising at numerous sites, including breast, ovary, lung, stomach,
10 and salivary gland (Hynes, N. E. & Stern, D. F. (1994) *Biochim. Biophys. Acta* **1198**, 165-
184). Increased expression of ErbB-2 leads to constitutive activation of its intrinsic tyrosine
kinase, and has been shown to cause the transformation of cultured cells. Numerous
15 clinical studies have shown that patients bearing tumors with elevated ErbB-2 expression
levels have a poorer prognosis (Hynes, N. E. & Stern, D. F. (1994) *Biochim. Biophys. Acta*
1198, 165-184). In addition to its involvement in human cancer, *erbB-2* plays important
biological roles, both in the adult and during embryonal development of mammals (Hynes,
20 N. E. & Stern, D. F. (1994) *Biochim. Biophys. Acta* **1198**, 165-184, Altioik, N., Bessereau,
J.-L. & Changeux, J.-P. (1995) *EMBO J.* **14**, 4258-4266, Lee, K.-F., Simon, H., Chen, H.,
Bates, B., Hung, M.-C. & Hauser, C. (1995) *Nature* **378**, 394-398).

The *erbB-2* promoter therefore represents an interesting test case for the
25 development of artificial transcriptional regulators. This promoter has been characterized in
detail and has been shown to be relatively complex, containing both a TATA-dependent and
a TATA-independent transcriptional initiation site (Ishii, S., Imamoto, F., Yamanashi, Y.,
Toyoshima, K. & Yamamoto, T. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 4374-4378).
30 Whereas early studies showed that polydactyl proteins could act as transcriptional
regulators that specifically activate or repress transcription, these proteins bound upstream
of an artificial promoter to six tandem repeats of the proteins binding site (Liu, Q., Segal, D.
J., Ghiara, J. B. & Barbas III, C. F. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 5525-5530).
35 Furthermore, this study utilized polydactyl proteins that were not modified in their binding
specificity. Herein, we tested the efficacy of polydactyl proteins assembled from predefined
building blocks to bind a single site in the native *erbB-2* promoter. Described above is the
generation and characterization of a family of zinc finger domains that bind each of the 16
40 5'-(GNN)-3' DNA triplets. One reason we focused on the production of this family of
recognition domains is that promoter regions of most organisms are relatively GC rich in
their base content. Thus, if proteins recognizing 5'-(GNN)_x-3' sites could be readily
assembled from this set of defined zinc finger domains, many genes could be rapidly and
45 specifically targeted for regulation. A protein containing six zinc finger domains and
recognizing 18 bp of DNA should be sufficient to define a single address within all known
genomes. Examination of the *erbB-2* promoter region revealed two 5'-(GNN)₆-3' sites and
50 one 5'-(GNN)₉-3' site. One of these sites, identified here as e2c, falls within the 5'-

5 untranscribed region of the *erbB-2* gene and was chosen as the target site for the generation
of a gene-specific transcriptional switch. A BLAST sequence similarity search of the
10 GenBank data base confirmed that this sequence is unique to *erbB-2*. The position of the
e2c target sequence, downstream and in the vicinity of the two major transcription initiation
sites, allowed for the examination of repression through inhibition of either transcription
15 initiation or elongation. An interesting feature of the e2c target site is that it is found within a
short stretch of sequence that is conserved between human, rat, and mouse *erbB-2* genes
(White, M. R.-A. & Hung, M.-C. (1992) *Oncogene* 7, 677-683). Thus, targeting of this site
would allow for the study of this strategy in animal models prior to its application to human
20 disease.

For generating polydactyl proteins with desired DNA-binding specificity, the present
studies have focused on the assembly of predefined zinc finger domains, which contrasts
the sequential selection strategy proposed by Greisman and Pabo (Greisman, H. A. &
25 Pabo, C. O. (1997) *Science* 275, 657-661). Such a strategy would require the sequential
generation and selection of six zinc finger libraries for each required protein, making this
experimental approach inaccessible to most laboratories and extremely time consuming to
all. Further, since it is difficult to apply specific negative selection against binding alternative
30 sequences in this strategy, proteins may result that are relatively unspecific as was recently
reported (Kim, J.-S. & Pabo, C. O. (1997) *J. Biol. Chem.* 272, 29795-29800).

The general utility of two different strategies for generating three-finger proteins
recognizing 9 bp of DNA sequence was investigated. Each strategy was based on the
35 modular nature of the zinc finger domain, and takes advantage of a family of zinc finger
domains recognizing triplets of the 5'-GNN-3'. Two three-finger proteins recognizing
half-sites (HS) 1 and 2 of the 5'-(GNN)₃-3' *erbB-2* target site e2c were generated in the first
strategy by fusing the pre-defined finger 2 (F2) domain variants together using a PCR
40 assembly strategy. To examine the generality of this approach, three additional three-finger
proteins recognizing sequences of the 5'-(GNN)₃-3' type, were prepared using the same
approach. Purified zinc finger proteins were prepared as fusions with the maltose binding
protein (MBP). ELISA analysis revealed that serially connected F2 proteins were able to act
45 in concert to specifically recognize the desired 9-bp DNA target sequences. Each of the 5
proteins shown was able to discriminate between target and non-target 5'-(GNN)₃-3'
sequence.

5

10

15

20

25

30

35

40

45

50

55

The affinity of each of the proteins for its target was determined by electrophoretic mobility-shift assays. These studies demonstrated that the zinc finger peptides have affinities comparable to Zif268 and other natural transcription factors with K_d values that ranged from 3 to 70 nM. Here the K_d of Zif268 for its operator to be 10 nM. It must be noted that, for reasons that remain to be explained, one group has reported K_d values for the natural Zif268 protein that range from 6 nM to 10 pM, a 600-fold variation (Pavletich, N. P. & Pabo, C. O. (1991) *Science* **252**, 809-17., Greisman, H. A. & Pabo, C. O. (1997) *Science* **275**, 657-661). Most studies have reported the K_d of the Zif268-DNA interaction to be from 3 to 10 nM, Choo, Y. & Klug, A. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 11163-11167, Hamilton, T. B., Borel, F. & Romaniuk, P. J. (1998) *Biochemistry* **37**, 2051-2058). Thus, in order to compare the results reported here with those reported elsewhere, the relative K_d s should be compared, $(\text{Mutant } K_d)/(\text{Zif268 } K_d)$, where both values are derived from the same report. The present data compare favorably to other studies of novel three-finger proteins prepared using phage display where affinities 10- to 200-fold weaker than Zif268 were reported (Greisman, H. A. & Pabo, C. O. (1997) *Science* **275**, 657-661, Choo, Y., Sanchez-Garcia, I. & Klug, A. (1994) *Nature* **372**, 642-5).

As an alternative to the serial connection of F2 domain variants, in the second strategy, three-finger proteins specific for the two e2c 5'-(GNN)₃-3' halfsites were produced by "helix grafting". The framework residues of the zinc finger domains, those residues that support the presentation of the recognition helix, vary between proteins. We anticipated that the framework residues may play a role in affinity and specificity. For helix grafting, amino acid positions -2 to 6 of the DNA recognition helices were either grafted into a Zif268 (Pavletich, N. P. & Pabo, C. O. (1991) *Science* **252**, 809-17) or an Sp1C framework (Desjarlais, J. R. & Berg, J. M. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 2256-60). The Sp1C protein is a designed consensus protein shown to have enhanced stability towards chelating agents. The proteins were expressed from DNA templates prepared by a rapid PCR-based gene assembly strategy. In each case, ELISA analysis of MBP fusion proteins showed that the DNA binding specificities and affinities observed with the F2 framework constructs were retained.

As discussed above, the recognition of 9 bp of DNA sequence is not sufficient to specify a unique site within a complex genome. In contrast, a six-finger protein recognizing 18 bp of contiguous DNA sequence could define a single site in the human genome, thus fulfilling an important prerequisite for the generation of a gene-specific transcriptional switch. Six-finger proteins binding the *erbB-2* target sequence e2c were generated from three-finger

5 constructs by simple restriction enzyme digestion and cloning with F2, Zif268, and Sp1C
framework template DNAs. ELISA analysis of purified MBP fusion proteins showed that
10 each of the six-finger proteins was able to recognize the specific target sequence, with little
cross reactivity to non-target 5'-(GNN)₆-3' sites or a tandem repeat of the Zif268 target site.

The affinity of each protein for the e2c DNA target site was determined by gel-shift
15 analysis. A modest K_d value of 25 nM was observed with the E2C(F2) six-finger protein
constructed from the F2 framework, a value that is only 2 to 3 times better than its
constituent three-finger proteins. In our previous studies of six-finger proteins, we observed
approximately 70-fold enhanced affinity of the six-finger proteins for their DNA ligand as
20 compared to their three-finger constituents (Liu, Q., Segal, D. J., Ghiara, J. B. & Barbas III,
C. F. (1997) *Proc. Natl. Acad. Sci. USA* 94, 5525-5530). The absence of a substantial
increase in the affinity of the E2C(F2) peptide suggested that serial connection of F2
domains is not optimal. It is possible that the periodicity of the F2 domains of the six-finger
25 protein does not match that of the DNA over this extended sequence, and that a significant
fraction of the binding energy of this protein is spent in unwinding DNA (Shi, Y. & Berg, J.
M. (1996) *Biochemistry* 35, 3845-8). In contrast to the F2 domain protein, the E2C(Zif) and
E2C(Sp1) six-finger proteins displayed 40- to 70-fold increased affinity as compared to their
30 original three-finger protein constituents, with K_d values of 1.6nM and 0.5nM, respectively.
Significantly, both three-finger components of these proteins were involved in binding, since
mutation of either half-site led to a roughly 100-fold decrease in affinity. The preponderance
of known transcription factors bind their specific DNA ligands with nanomolar affinity,
35 suggesting that the control of gene expression is governed by protein/DNA complexes of
unexceptional life times. Thus, zinc finger proteins of increased affinity should not be
required and could be disadvantageous, especially if binding to non-specific DNA is also
increased.

40 The zinc finger domain is generally considered to be modular in nature, with each finger
recognizing a 3-bp subsite (Pavletich, N. P. & Pabo, C. O. (1991) *Science* 252, 809-17).
This is supported by our ability to recombine zinc finger domains in any desired sequence,
yielding polydactyl proteins recognizing extended sequences of the structure 5'-(GNN)_x-3'.
45 However, it should be noted that at least in some cases, zinc finger domains appear to
specify overlapping 4 bp sites rather than individual 3 bp sites. In Zif268, residues in
addition to those found at helix positions -1, 3, and 6 are involved in contacting DNA (Elrod-
50 Erickson, M., Rould, M. A., Neklodova, L. & Pabo, C. O. (1996) *Structure* 4, 1171-1180).

Specifically, an aspartate in helix position 2 of F2 plays several roles in recognition and makes a variety of contacts. The carboxylate of the aspartate side chain hydrogen bonds with arginine at position -1, stabilizing its interaction with the 3'-guanine of its target site. This aspartate also participates in water-mediated contacts with the guanine's complementary cytosine. In addition, this carboxylate is observed to make a direct contact to the N4 of the cytosine base on the opposite strand of the 5'-guanine base of the finger 1 binding site. It is this interaction which is the chemical basis for target site overlap. Indeed, when the Zif268 F2 libraries were selected against the four 5'-GCG GNG GCG-3' sequences, both an arginine at position -1 and an aspartate at position 2 were obtained, analogous to the residues in native Zif268. Since the e2c target sequence (5'-GGG GCC GGA GCC GCA GTG-3') (SEQ ID NO: 112) is followed by an A rather than a G, a potential target site overlap problem was anticipated with finger 1 of an e2c-specific six-finger protein. However, in both the Zif- and Sp1C-framework six-finger proteins, the GTG-specific finger 1 containing an aspartate at position 2 appears to recognize the sequences 5'-GTGA-3' and 5'-GTGG-3' equally well, as indicated by their very similar affinities to target sites e2c-a and e2c-g.

A polynucleotide or composition of this invention as set forth above, can be operatively linked to one or more transcription modulating factors. Modulating factors such as transcription activators or transcription suppressors or repressors are well known in the art. Means for operatively linking polypeptides to such factors are also well known in the art. Exemplary and preferred such factors and their use to modulate gene expression are discussed in detail hereinafter.

II Uses

In one embodiment, a method of the invention includes a process for modulating (inhibiting or suppressing) the function of a nucleotide sequence comprising a zinc finger-nucleotide binding motif which comprises contacting the zinc finger-nucleotide binding motif with an effective amount of a zinc finger-nucleotide binding polypeptide that binds to the motif. In the case where the nucleotide sequence is a promoter, the method includes inhibiting the transcriptional transactivation of a promoter containing a zinc finger-DNA binding motif. The term "inhibiting" refers to the suppression of the level of activation of transcription of a structural gene operably linked to a promoter, containing a zinc finger-nucleotide binding motif, for example. In addition, the zinc finger-nucleotide binding polypeptide derivative may bind a motif within a structural gene or within an RNA sequence.

5

10

15

20

25

The term "effective amount" includes that amount which results in the deactivation of a previously activated promoter or that amount which results in the inactivation of a promoter containing a zinc finger-nucleotide binding motif, or that amount which blocks transcription of a structural gene or translation of RNA. The amount of zinc finger derived-nucleotide binding polypeptide required is that amount necessary to either displace a native zinc finger-nucleotide binding protein in an existing protein/promoter complex, or that amount necessary to compete with the native zinc finger-nucleotide binding protein to form a complex with the promoter itself. Similarly, the amount required to block a structural gene or RNA is that amount which binds to and blocks RNA polymerase from reading through on the gene or that amount which inhibits translation, respectively. Preferably, the method is performed intracellularly. By functionally inactivating a promoter or structural gene, transcription or translation is suppressed. Delivery of an effective amount of the inhibitory protein for binding to or "contacting" the cellular nucleotide sequence containing the zinc finger-nucleotide binding protein motif, can be accomplished by one of the mechanisms described herein, such as by retroviral vectors or liposomes, or other methods well known in the art.

30

35

40

The term "modulating" refers to the suppression, enhancement or induction of a function. For example, the zinc finger-nucleotide binding polypeptide of the invention may modulate a promoter sequence by binding to a motif within the promoter, thereby enhancing or suppressing transcription of a gene operatively linked to the promoter nucleotide sequence. Alternatively, modulation may include inhibition of transcription of a gene where the zinc finger-nucleotide binding polypeptide binds to the structural gene and blocks DNA dependent RNA polymerase from reading through the gene, thus inhibiting transcription of the gene. The structural gene may be a normal cellular gene or an oncogene, for example. Alternatively, modulation may include inhibition of translation of a transcript.

45

The promoter region of a gene includes the regulatory elements that typically lie 5' to a structural gene. If a gene is to be activated, proteins known as transcription factors attach to the promoter region of the gene. This assembly resembles an "on switch" by enabling an enzyme to transcribe a second genetic segment from DNA to RNA. In most cases the resulting RNA molecule serves as a template for synthesis of a specific protein; sometimes RNA itself is the final product.

50

The promoter region may be a normal cellular promoter or, for example, an onco-promoter. An onco-promoter is generally a virus-derived promoter. For example, the long

55

5

10

15

20

25

30

35

40

45

50

55

terminal repeat (LTR) of retroviruses is a promoter region which may be a target for a zinc finger binding polypeptide variant of the invention. Promoters from members of the Lentivirusgroup, which include such pathogens as human T-cell lymphotropic virus (HTLV) 1 and 2, or human immunodeficiency virus (HIV) 1 or 2, are examples of viral promoter regions which may be targeted for transcriptional modulation by a zinc finger binding polypeptide of the invention.

In order to test the concept of using zinc finger proteins as gene-specific transcriptional regulators, the E2C(Sp1) six-finger protein was fused to a number of effector domains. Transcriptional repressors were generated by attaching either of three human-derived repressor domains to the zinc finger protein. The first repressor protein was prepared using the ERF repressor domain (ERD) (Sgouras, D. N., Athanasiou, M. A., Beal, G. J., Jr., Fisher, R. J., Blair, D. G. & Mavrothalassitis, G. J. (1995) *EMBO J.* **14**, 4781-4793), defined by amino acids 473 to 530 of the *ets2* repressor factor (ERF). This domain mediates the antagonistic effect of ERF on the activity of transcription factors of the *ets* family. A synthetic repressor was constructed by fusion of this domain to the C-terminus of the zinc finger protein. The second repressor protein was prepared using the Krüppel-associated box (KRAB) domain (Margolin, J. F., Friedman, J. R., Meyer, W., K.-H., Vissing, H., Thiesen, H.-J. & Rauscher III, F. J. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 4509-4513). This repressor domain is commonly found at the N-terminus of zinc finger proteins and presumably exerts its repressive activity on TATA-dependent transcription in a distance- and orientation-independent manner (Pengue, G. & Lania, L. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 1015-1020), by interacting with the RING finger protein KAP-1 (Friedman, J. R., Fredericks, W. J., Jensen, D. E., Speicher, D. W., Huang, X.-P., Neilson, E. G. & Rauscher III, F. J. (1996) *Genes & Dev.* **10**, 2067-2078). We utilized the KRAB domain found between amino acids 1 and 97 of the zinc finger protein KOX1 (Margolin, J. F., Friedman, J. R., Meyer, W., K.-H., Vissing, H., Thiesen, H.-J. & Rauscher III, F. J. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 4509-4513). In this case an N-terminal fusion with the six-finger protein was constructed. Finally, to explore the utility of histone deacetylation for repression, amino acids 1 to 36 of the Mad mSIN3 interaction domain (SID) were fused to the N-terminus of the zinc finger protein (Ayer, D. E., Laherty, C. D., Lawrence, Q. A., Armstrong, A. P. & Eisenman, R. N. (1996) *Mol. Cell. Biol.* **16**, 5772-5781). This small domain is found at the N-terminus of the transcription factor Mad and is responsible for mediating its transcriptional repression by interacting with mSIN3, which in turn interacts the co-repressor N-CoR and

5 with the histone deacetylase mRPD1 (Heinzel, T., Lavinsky, R. M., Mullen, T.-M.,
Söderström, M., Laherty, C. D., Torchia, J., Yang, W.-M., Brard, G., Ngo, S. D. & al., e.
10 (1997) *Nature* **387**, 43-46). To examine gene-specific activation, transcriptional activators
were generated by fusing the zinc finger protein to amino acids 413 to 489 of the herpes
simplex virus VP16 protein (Sadowski, I., Ma, J., Triezenberg, S. & Ptashne, M. (1988)
15 *Nature* **335**, 563-564), or to an artificial tetrameric repeat of VP16's minimal activation
domain, DALDDFDLML (SEQ ID NO:113) (Seipel, K., Georgiev, O. & Schaffner, W.
(1992) *EMBO J.* **11**, 4961-4968), termed VP64.

Reporter constructs containing fragments of the *erbB-2* promoter coupled to a luciferase
reporter gene were generated to test the specific activities of our designed transcriptional
20 regulators. The target reporter plasmid contained nucleotides -758 to -1 with respect to the
ATG initiation codon, whereas the control reporter plasmid contained nucleotides -1571 to -
24, thus lacking all but one nucleotide of the E2C binding site encompassed in positions -24
25 to -7. Both promoter fragments displayed similar activities when transfected transiently into
HeLa cells, in agreement with previous observations (Hudson, L. G., Ertl, A. P. & Gill, G. N.
(1990) *J. Biol. Chem.* **265**, 4389-4393). To test the effect of zinc finger-repressor domain
fusion constructs on *erbB-2* promoter activity, HeLa cells were transiently co-transfected
30 with each of the zinc finger expression vectors and the luciferase reporter constructs (Fig.
5A). Significant repression was observed with each construct. The ERD and SID fusion
proteins produced approximately 50% and 80% repression, respectively. The most potent
repressor was the KRAB fusion protein. This protein caused complete repression of *erbB-2*
35 promoter activity. The observed residual activity was at the background level of the
promoter-less pGL3 reporter. In contrast, none of the proteins caused significant repression
of the control *erbB-2* reporter construct lacking the E2C target site, demonstrating that
repression is indeed mediated by specific binding of the E2C(Sp1) protein to its target site.
40 Expression of a zinc finger protein lacking any effector domain resulted in weak repression,
approximately 30%, indicating that most of the repression observed with the SID and KRAB
constructs is caused by their effector domains, rather than by DNA-binding alone. This
observation strongly suggests that the mechanism of repression is active inhibition of
45 transcription initiation rather than of elongation. Once initiation of transcription by RNA
polymerase II has occurred, the zinc finger protein appears to be readily displaced from the
DNA by the action of the polymerase.

5
10
15
The utility of gene-specific polydactyl proteins to mediate activation of transcription was investigated using the same two reporter constructs. The VP16 fusion protein was found to stimulate transcription approximately 5-fold, whereas the VP64 fusion protein produced a 27-fold activation. This dramatic stimulation of promoter activity caused by a single VP16-based transcriptional activator is exceptional in view of the fact that the zinc finger protein binds in the transcribed region of the gene. This again demonstrates that mere binding of a zinc finger protein, even with one with subnanomolar affinity, in the path of RNA polymerase II need not necessarily negatively affect gene expression.

20
25
30
35
40
45
50
55
The data herein show that zinc finger proteins capable of binding novel 9- and 18-bp DNA target sites can be rapidly prepared using pre-defined domains recognizing 5'-GNN-3' sites. This information is sufficient for the preparation of 16⁸ or 17 million novel six-finger proteins each capable of binding 18 bp of DNA sequence. This rapid methodology for the construction of novel zinc finger proteins has advantages over the sequential generation and selection of zinc finger domains proposed by others (Greisman, H. A. & Pabo, C. O. (1997) *Science* **275**, 657-661) and takes advantage of structural information that suggests that the potential for the target overlap problem as defined above might be avoided in proteins targeting 5'-GNN-3' sites. Using the complex and well studied *erbB-2* promoter and live human cells, the data demonstrate that these proteins, when provided with the appropriate effector domain, can be used to provoke or activate expression and to produce graded levels of repression down to the level of the background in these experiments. These studies suggest that the KRAB domain is significantly more potent as a transcriptional repressor than ERD or SID domains, and that it is able to inhibit both the TATA-dependent and the TATA-independent transcriptional initiation of this promoter. These repressor domains have not previously been directly compared. The present strategy of using predefined zinc finger domains to construct polydactyl proteins coupled to effector domains has significant advantages over strategies that attempt to only repress transcription by competing or interfering with proteins involved in the transcription complex (Kim, J.-S. & Pabo, C. O. (1997) *J. Biol. Chem.* **272**, 29795-29800, Kim, J.-S., Kim, J., Cepek, K. L., Sharp, P. A. & Pabo, C. O. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 3616-3620). Utilization of effector domains that have the potential to act over a distance should allow the application of these gene-switches to the regulation of uncharacterized genes and promoters. Since these transcriptional regulators might be prepared using our PCR-assembly strategy in a high-throughput fashion, we believe it is appropriate to comment on their potential practical applications. Novel DNA binding proteins generated in this manner

5 should have potential utility in DNA-based diagnostic applications. For the study of gene
function, we believe that the ability to both activate and repress the transcription of genes,
10 at graded levels if necessary, may assist in assigning gene function. Since these proteins
exert their control by acting in *trans*, functional gene knockout or activation might be
produced in heterozygous transgenic animals. This would drastically reduce the time
required to produce a gene knockout in a whole animal and would extend the range of
15 organisms to which knockout technology might be applied. These proteins might also be
used in gene therapy applications to inhibit the production of viral gene products or to
activate genes involved in fighting disease. Significantly, the ease with which these proteins
can be prepared will facilitate the testing of these ideas by the scientific community.

20 The Examples that follow illustrate preferred embodiments of the present invention
and are not limiting of the specification or claims in any way.

25 **EXAMPLE 1: Selection by phage display.**

Construction of zinc-finger libraries by PCR overlap extension was essentially as
previously described (Shi, Y. & Berg, J. M. (1996) *Biochemistry* 35, 3845-8). Growth and
precipitation of phage were as previously described (Pengue, G. & Lania, L. (1996) *Proc.*
30 *Natl. Acad. Sci. USA* 93, 1015-1020, Friedman, J. R., Fredericks, W. J., Jensen, D. E.,
Spelcher, D. W., Huang, X.-P., Neilson, E. G. & Rauscher III, F. J. (1996) *Genes & Dev.* 10,
2067-2078), except that ER2537 cells (New England Biolabs) were used to propagate the
phage and 90 μ M ZnCl₂ was added to the growth media. Precipitated phage were
35 resuspended in Zinc Buffer A (ZBA; 10 mM Tris, pH7.5/90 mM KCl, 1 mM MgCl₂, 90 μ M
ZnCl₂)/1% BSA /5 mM DTT. Binding reactions (500 μ l: ZBA/5 mM DTT/1% Blotto
(BioRad)/competitor oligonucleotides/4 μ g sheared herring sperm DNA (Sigma)/100 μ l
40 filtered phage ($\approx 10^{13}$ colony forming units)) were incubated for 30 minutes at room
temperature, prior to the addition of 72nM biotinylated hairpin target oligonucleotide.
Incubation continued for 3.5 hours with constant gentle mixing. Streptavidin-coated
magnetic beads (50 μ l; Dynal) were washed twice with 500 μ l ZBA/1% BSA, then blocked
45 with 500 μ l ZBA/5% Blotto/ antibody-displaying (irrelevant) phage ($\approx 10^{12}$ colony forming
units) for ≈ 4 hours at room temperature. At the end of the binding period, the blocking
solution was replaced by the binding reaction and incubated 1 hour at room temperature.
50 The beads were washed 10 times over a 1 hour period with 500 μ l ZBA/5 mM DTT/2%

5 Tween 20, then once without Tween 20. Bound phage were eluted 30 minutes with 10 $\mu\text{g}/\mu\text{l}$ trypsin.

10 Hairpin target oligonucleotides had the sequence 5'-Biotin-GGACGCN'N'N'CGCGGGTTTCCCGCGNNGCGTCC-3' (SEQ ID NO:114), where NNN was the 3-nucleotide finger 2-target sequence and N'N'N' its complement. A similar nonbiotinylated oligonucleotide, in which the target sequence was TGG (compTGG), was
15 included at 7.2nM in every round of selection to select against contaminating parental phage. Two pools of nonbiotinylated oligonucleotides were also used as competitors: one containing all 64 possible 3-nucleotide targets sequences (compNNN), the other containing all the GNN target sequences except for the current selection target (compGNN). These
20 pools were typically used as follows: round 1, no compNNN or compGNN; round 2, 7.2nM compGNN; round 3, 10.8nM compGNN; round 4, 1.8 μM compNNN, 25nM compGNN; round 5, 2.7 μM compNNN, 90nM compGNN; round 6, 2.7 μM compNNN, 250nM compGNN; round
25 7, 3.6 μM compNNN, 250nM compGNN.

EXAMPLE 2: Multi-target specificity assays.

The fragment of pComb3H (Pengue, G. & Lania, L. (1996) *Proc. Natl. Acad. Sci. USA* 93, 1015-1020, Heinzel, T., Lavinsky, R. M., Mullen, T.-M., Söderström, M., Laherty, C. D.,
30 Torchia, J., Yang, W.-M., Brard, G., Ngo, S. D. & al., e. (1997) *Nature* 387, 43-46) phagemid RF DNA containing the zinc-finger coding sequence was subcloned into a modified pMAL-c2 (New England Biolabs) bacterial expression vector and transformed into
35 XL1-Blue (Stratagene). Freeze/thaw extracts containing the overexpressed maltose binding protein-zinc finger fusion proteins were prepared from IPTG-induced cultures using the Protein Fusion and Purification System (New England Biolabs). In 96-well ELISA plates, 0.2 μg of streptavidin (Pierce) was applied to each well for 1 hour at 37°C, then washed
40 twice with water. Biotinylated target oligonucleotide (0.025 μg) was applied similarly. ZBA/3% BSA was applied for blocking, but the well were not washed after incubation. All subsequent incubations were at room temperature. Eight 2-fold serial
45 dilutions of the extracts were applied in 1x binding buffer (ZBA/1% BSA/5 mM DTT/0.12 $\mu\text{g}/\mu\text{l}$ sheared herring sperm DNA). The samples were incubated 1 hour, followed by 10 washes with water. Mouse anti-maltose binding protein mAb (Sigma) in ZBA/1% BSA was applied to the wells for 30 minutes, followed by 10 washes with water. Goat anti -mouse IgG
50 mAb conjugated to alkaline phosphatase (Sigma) was applied to the wells for 30 minutes,

5 followed by 10 washes with water. Alkaline phosphatase substrate (Sigma) was applied, and the OD₄₀₅ was quantitated with SOFTmax 2.35 (Molecular Devices).

10 **EXAMPLE 3: Gel mobility shift assays.**

Fusion proteins were purified to >90% homogeneity using the Protein Fusion and Purification System (New England Biolabs), except that ZBA/5 mM DTT was used as the column buffer. Protein purity and concentration were determined from Coomassie blue-stained 15% SDS-PAGE gels by comparison to BSA standards. Target oligonucleotides were labeled at their 5' or 3' ends with [³²P] and gel purified. Eleven 3-fold serial dilutions of protein were incubated in 20 µl binding reactions (1xBinding Buffer/10% glycerol/1 pM target oligonucleotide) for three hours at room temperature, then resolved on a 5% polyacrylamide gel in 0.5xTBE buffer. Quantitation of dried gels was performed using a PhosphorImager and ImageQuant software (Molecular Dynamics), and the K_D was determined by scatchard analysis.

25 **EXAMPLE 4: Generation of polydactyl proteins with desired DNA binding specificity.**

The studies reported here use the finger 2 (F2) variants pmGAC, pmGAG, pGCA, pGCC, pmGGA, pmGGC, pmGGG, and pGTG defined in the accompanying manuscript (Hudson, L. G., Ertl, A. P. & Gill, G. N. (1990) *J. Biol. Chem.* **265**, 4389-4393). To generate DNAs encoding three-finger proteins, F2 coding regions were PCR amplified from selected or designed F2 variants and assembled by PCR overlap extension. Alternatively, DNAs encoding three-finger proteins with a Zif268 or Sp1C framework were synthesized from 8 or 6 overlapping oligonucleotides, respectively. Sp1C framework constructs, used for all reporter assays described in this report, were generated as follows. In the case of E2C-HS1(Sp1), 0.4 pmole each of oligonucleotides SPE2-3 (5'-GCG AGC AAG GTC GCG GCA GTC ACT AAA AGA TTT GCC GCA CTC TGG GCA TTT ATA CGG TTT TTC ACC-3') (SEQ ID NO:115) and SPE2-4 (5'-GTG ACT GCC GCG ACC TTG CTC GCC ATC AAC GCA CTC ATA CTG GCG AGA AGC CAT ACA AAT GTC CAG AAT GTG GC-3') (SEQ ID NO:116) were mixed with 40 pmole each of oligonucleotides SPE2-2 (5'-GGT AAG TCC TTC TCT CAG AGC TCT CAC CTG GTG CGC CAC CAG CGT ACC CAC ACG GGT GAA AAA CCG TAT AAA TGC CCA GAG-3') (SEQ ID NO:117) and SPE2-5 (5'-ACG CAC CAG CTT GTC AGA GCG GCT GAA AGA CTT GCC ACA TTC TGG ACA TTT GTA TGG C-3') (SEQ ID NO:118) in a standard PCR mixture and cycled 25 times (30 seconds at 94°C, 30

seconds at 60°C, 30 seconds at 72°C). An aliquot of this pre-assembly reaction was then amplified with 40 pmole each of the primers SPE2-1 (5'-GAG GAG GAG GAG GTG GCC CAG GCG GCC CTC GAG CCC GGG GAG AAG CCC TAT GCT TGT CCG GAA TGT GGT AAG TCC TTC TCT CAG AGC-3') (SEQ ID NO:119) and SPE2-6 (5'-GAG GAG GAG GAG CTG GCC GGC CTG GCC ACT AGT TTT TTT ACC GGT GTG AGT ACG TTG GTG ACG CAC CAG CTT GTC AGA GCG-3') (SEQ ID NO:120) using the same cycling conditions. The E2C-HS2(Sp1) DNA was generated in the same way, using an analogous set of oligonucleotides differing only in the recognition helix coding regions. All assembled three-finger coding regions were digested with the restriction endonuclease *Sfi*I and cloned into pMal-CSS, a derivative of the bacterial expression vector pMal-C2 (New England Biolabs). DNAs encoding six-finger proteins with each of the different frameworks were assembled in pMal-CSS using *Xma*I and *Bsr*FI restriction sites included in the sequences flanking the three-finger coding regions. Each of the zinc finger proteins was expressed in the *E. coli* strain XL1-blue and binding properties were investigated by ELISA and gel shift analysis as described in the accompanying manuscript (Hudson, L. G., Ertl, A. P. & Gill, G. N. (1990) *J. Biol. Chem.* 265, 4389-4393).

EXAMPLE 5: Construction of zinc finger-effector domain fusion proteins.

For the construction of zinc finger-effector domain fusion proteins, DNAs encoding amino acids 473 to 530 of the *gts* repressor factor (ERF) repressor domain (ERD) (Sgouras, D. N., Athanasiou, M. A., Beal, G. J., Jr., Fisher, R. J., Blair, D. G. & Mavrothalassitis, G. J. (1995) *EMBO J.* 14, 4781-4793), amino acids 1 to 97 of the KRAB domain of KOX1 (Margolin, J. F., Friedman, J. R., Meyer, W., K.-H., Vissing, H., Thiesen, H.-J. & Rauscher III, F. J. (1994) *Proc. Natl. Acad. Sci. USA* 91, 4509-4513), or amino acids 1 to 36 of the Mad mSIN3 interaction domain (SID) (Ayer, D. E., Laherty, C. D., Lawrence, Q. A., Armstrong, A. P. & Eisenman, R. N. (1996) *Mol. Cell. Biol.* 16, 5772-5781) were assembled from overlapping oligonucleotides using Taq DNA polymerase. The coding region for amino acids 413 to 489 of the VP16 transcriptional activation domain (Sadowski, I., Ma, J., Triezenberg, S. & Ptashne, M. (1988) *Nature* 335, 563-564) was PCR amplified from pcDNA3/C7-C7-VP16 (10). The VP64 DNA, encoding a tetrameric repeat of VP16's minimal activation domain, comprising amino acids 437 to 447 (Seipel, K., Georgiev, O. & Schaffner, W. (1992) *EMBO J.* 11, 4961-4968), was generated from two pairs of complementary oligonucleotides. The resulting fragments were fused to zinc finger coding

5 regions by standard cloning procedures, such that each resulting construct contained an
internal SV40 nuclear localization signal, as well as a C-terminal HA decapeptide tag.
10 Fusion constructs were cloned in the eucaryotic expression vector pcDNA3 (Invitrogen).

EXAMPLE 6: Construction of luciferase reporter plasmids.

15 An *erbB-2* promoter fragment comprising nucleotides -758 to -1, relative to the ATG
initiation codon, was PCR amplified from human bone marrow genomic DNA with the
TaqExpand DNA polymerase mix (Boehringer Mannheim) and cloned into pGL3basic
(Promega), upstream of the firefly luciferase gene. A human *erbB-2* promoter fragment
20 encompassing nucleotides -1571 to -24, was excised from pSVOALΔ5'/*erbB-2*(N-N)
(Hudson, L. G., Ertl, A. P. & Gill, G. N. (1990) *J. Biol. Chem.* 265, 4389-4393) by *Hind3*
digestion and subcloned into pGL3basic, upstream of the firefly luciferase gene.

EXAMPLE 7: Luciferase assays.

25 For all transfections, HeLa cells were used at a confluency of 40-60%. Typically, cells
were transfected with 400 ng reporter plasmid (pGL3-promoter constructs or, as negative
control, pGL3basic), 50 ng effector plasmid (zinc finger constructs in pcDNA3 or, as
30 negative control, empty pcDNA3), and 200 ng internal standard plasmid (p β Act- β Gal) in a
well of a 6 well dish using the lipofectamine reagent (Gibco BRL). Cell extracts were
prepared approximately 48 hours after transfection. Luciferase activity was measured with
luciferase assay reagent (Promega), β Gal activity with Galacto-Light (Tropix), in a
35 MicroLumat LB96P luminometer (EG&G Berthold). Luciferase activity was normalized on
 β Gal activity.

EXAMPLE 8: Regulation of the *erbB-2* gene in HeLa cells.

40 The *erbB-2* gene was targeted for imposed regulation. The *erbB-2* gene is frequently
overexpressed in human cancers, particularly breast and ovarian, and elevated *ErbB-2*
levels correlate with a poor prognosis (N. E. Hynes and D. F. Sten, *Biochim. Biophys. Acta*
1198, 165 (1994)). To regulate the native *erbB-2* gene, a synthetic repressor protein,
45 designated E2C-KRAB, and a transactivator protein, designated E2C-VP64, were utilized
(R. R. Beerli, D. J. Segal, B. Dreier, C. F. Barbas, III, *Proc. Natl. Acad. Sci. USA* 95, 14628
(1998)). Both proteins contain the same designed zinc finger protein E2C that recognizes
50 the 18-bp DNA sequence 5'-GGG GCC GGA GCC GCA GTG-3' (SEQ ID NO:121) in the 5'-

5 untranscribed region of the proto-oncogene *erbB-2*. This DNA-binding protein was
constructed from 6 pre-defined and modular zinc finger domains (D. J. Segal, B. Dreier, R.
10 R. Beerli, C. F. Barbas, III, *Proc. Natl. Acad. Sci. USA* 96, 2758 (1999)). The repressor
protein contains the Kox-1 KRAB domain (J. F. Margolin *et al.*, *Proc. Natl. Acad. Sci. USA*
91, 4509 (1994)), whereas the transactivator VP64 contains a tetrameric repeat of the
minimal activation domain (K. Seipel, O. Georgiev, W. Schaffner, *EMBO J.* 11, 4961 (1992))
15 derived from the herpes simplex virus protein VP16.

A derivative of the human cervical carcinoma cell line HeLa, HeLa/tet-off, was utilized
(M. Gossen and H. Bujard, *Proc. Natl. Acad. Sci. USA* 89, 5547 (1992)). Since HeLa cells
20 are of epithelial origin they express *ErbB-2* and are well suited for studies of *erbB-2* gene
targeting. HeLa/tet-off cells produce the tetracycline-controlled transactivator, allowing
induction of a gene of interest under the control of a tetracycline response element (TRE)
by removal of tetracycline or its derivative doxycycline (Dox) from the growth medium. We
25 have used this system to place our transcription factors under chemical control. Thus, the
pRevTRE/E2C-SKD and pRevTRE/E2C-VP64 plasmids were constructed (The E2C(Sp1)-
KRAB and E2C(Sp1)-VP64 coding regions were PCR amplified from pcDNA3-based
expression plasmids (R. R. Beerli, D. J. Segal, B. Dreier, C. F. Barbas, III, *Proc. Natl. Acad.*
Sci. USA 95, 14628 (1998)) and subcloned into pRevTRE (Clontech) using BamH1 and
30 Cla1 restriction sites, and into pMX-IRES-GFP [X. Liu *et al.*, *Proc. Natl. Acad. Sci. USA* 94,
10669 (1997)] using BamH1 and Not1 restriction sites. Fidelity of the PCR amplification was
confirmed by sequencing, transfected into HeLa/tet-off cells, and 20 stable clones each
35 were isolated and analyzed for Dox-dependent target gene regulation (The pRevTRE/E2C-
KRAB and pRevTRE/E2C-VP64 constructs were transfected into the HeLa/tet-off cell line
(M. Gossen and H. Bujard, *Proc. Natl. Acad. Sci. USA* 89, 5547 (1992)) using
Lipofectamine Plus reagent (Gibco BRL). After two weeks of selection in hygromycin-
40 containing medium, in the presence of 2 µg/ml Dox, stable clones were isolated and
analyzed for Dox-dependent regulation of *ErbB-2* expression. Western blots,
immunoprecipitations, Northern blots, and flow cytometric analyses were carried out
essentially as described [D. Graus-Porta, R. R. Beerli, N. E. Hynes, *Mol. Cell. Biol.* 15, 1182
45 (1995)]. As a read-out of *erbB-2* promoter activity, *ErbB-2* protein levels were initially
analyzed by Western blotting. A significant fraction of these clones showed regulation of
ErbB-2 expression upon removal of Dox for 4 days, i.e. downregulation of *ErbB-2* in E2C-
KRAB clones and upregulation in E2C-VP64 clones. *ErbB-2* protein levels were correlated
50

5 with altered levels of their specific mRNA, indicating that regulation of ErbB-2 expression
was a result of repression or activation of transcription. The additional ErbB-2 protein
10 expressed in E2C-VP64 clones was indistinguishable from naturally expressed protein and
biologically active, since epidermal growth factor (EGF) readily induced its tyrosine
phosphorylation. The ErbB-2 levels in the E2C-KRAB clone #27, in the absence of Dox,
were below the level of detection as was its EGF-induced tyrosine phosphorylation.
15 Therefore, ErbB-2 expression was also analyzed by flow cytometry, revealing no detectable
ErbB-2 expression in E2C-KRAB clone #27, in sharp contrast to the dramatic upregulation
(5.6 fold) of ErbB-2 in E2C-VP64 clone #18. Thus, the extent of erbB-2 gene regulation
ranged from total repression (E2C-KRAB clone #27) to almost 6-fold activation (E2C-VP64
20 clone #18). No significant effect on the expression of the related ErbB-1 protein was
observed, indicating that regulation of ErbB-2 expression was not a result of general down-
or up-regulation of transcription. In contrast to the efficacy of these transcription factors that
target 18 bps of DNA sequence using six zinc finger domains, transcriptional activators
25 prepared with three zinc finger domains that bind either of the 9-bp half-sites of the E2C
target sequence were unable to activate transcription of an erbB-2-luciferase reporter.
These results suggest that the increased specificity and affinity of six finger proteins may be
required to provide a dominant effect on gene regulation.
30

**EXAMPLE 9: Introduction of the coding regions of the E2C-KRAB and E2C-VP64
proteins into the retroviral vector pMX-IRES-GFP.**

35 In order to express the E2C-KRAB and E2C-VP64 proteins in several other cell lines,
their coding regions were introduced into the retroviral vector pMX-IRES-GFP. The
E2C(Sp1)-KRAB and E2C(Sp1)-VP64 coding regions were PCR amplified from pcDNA3-
based expression plasmids (R. R. Beerli, D. J. Segal, B. Dreier, C. F. Barbas, III, *Proc. Natl.*
40 *Acad. Sci. USA* 95, 14628 (1998)) and subcloned into pRevTRE (Clontech) using BamH1
and Cla1 restriction sites, and into pMX-IRES-GFP [X. Liu *et al.*, *Proc. Natl. Acad. Sci. USA*
94, 10669 (1997)] using BamH1 and Not1 restriction sites. Fidelity of the PCR amplification
was confirmed by sequencing. This vector expresses a single bicistronic message for the
45 translation of the zinc finger protein and, from an internal ribosome-entry site (IRES), the
green fluorescent protein (GFP). Since both coding regions share the same mRNA, their
expression is physically linked to one another and GFP expression is an indicator of zinc
finger expression. Virus prepared from these plasmids was then used to infect the human
50 carcinoma cell line A431 (pMX-IRES-GFP/E2C-KRAB and pMX-IRES-GFP/E2C-VP64
55

5 Plasmids were transiently transfected into the amphotropic packaging cell line Phoenix
Ampho using Lipofectamine Plus (Gibco BRL) and, two days later, culture supernatants
10 were used for infection of target cells in the presence of 8 µg/ml polybrene. Three days after
infection, cells were harvested for analysis). Three days after infection, ErbB-2 expression
was measured by flow cytometry. Significantly, about 59% of the E2C-KRAB virus treated
15 cells were essentially ErbB-2 negative, while in about 27% of the E2C-VP64 virus treated
cells ErbB-2 levels were increased. Plotting of GFP fluorescence vs. ErbB-2 fluorescence
revealed that there were two cell populations, one with normal ErbB-2 levels that was GFP
negative, and another with altered ErbB-2 levels that was GFP positive. Specificity of gene
20 targeting was investigated by measuring the expression levels of the related ErbB-1 and
ErbB-3 proteins. No significant alterations of these protein levels were detected, indicating
that erbB-2 gene targeting is specific and not a non-specific result of general alterations in
gene expression or overexpression of the effector domains. The lack of any appreciable
25 regulation of erbB-3 is particularly remarkable since its 5'-UTR contains the 18bp sequence
5'-GGa GCC GGA GCC GgA GTc-3' (SEQ ID NO:122), that presents only 3 mismatches to
E2C's designed target sequence (15bp identity - lowercase letters indicate differences) (M.
H. Kraus, W. Issing, T. Mikl, N. C. Popescu, S. A. Aaronson, *Proc. Natl. Acad. Sci. USA* 86,
30 9193 (1989)).

EXAMPLE 10: Regulation of the erbB-2 gene in non-human primate cells.

35 The zinc finger target sequence within erbB-2's 5'-UTR lays within a 28-bp sequence
stretch that is conserved in many species. To investigate regulation of erbB-2 gene
expression in non-human primate cells, COS-7 fibroblasts were infected with the bicistronic
E2C-KRAB retrovirus and analyzed by flow cytometry. As in human cells, expression of the
40 repressor protein as indicated by the GFP marker correlated well with a loss of ErbB-2
protein. Similarly, gene targeting in murine cells was evaluated by infection of NIH/3T3 cells
with E2C-KRAB and E2C-VP64 encoding retrovirus. ErbB-2 expression levels were then
monitored by Western blotting rather than flow cytometry, due to a lack of reactivity of the
45 mAb with the murine ErbB-2 extracellular domain. Here again, with E2C-KRAB a complete
transcriptional knockout upon correction for infected cells was observed. However, unlike in
human cell lines, E2C-VP64 induced ErbB-2 upregulation was rather modest in NIH/3T3
cells, approximately 1.8 fold upon correction for infection efficiency. A likely explanation for
50 this discrepancy lies in the different structures of the human and mouse promoters. The
mouse erbB-2 promoter, unlike the human, does not contain a TATA box (M. R. White and

5 M. C. Hung, *Oncogene* 7, 677 (1992)). Transcriptional activation by VP16 is, at least in part,
mediated by its interaction with TFIID, a multi-protein complex also containing the TATA-
10 binding protein (C. J. Ingles, M. Shales, W. D. Cress, S. J. Triezenberg, J. Greenblatt,
Nature 351, 588 (1991)). It is therefore plausible that the E2C-VP64 protein activates
transcription less effectively in the absence of a TATA box. These data suggest that while a
15 DNA binding site may be conserved with respect to sequence and relative position within a
target cell, effector domains may need to be optimized for maximal efficiency due to context
effects. Nevertheless, while their potencies may differ, the artificial transcription factors
described here are capable of imposing regulation of erbB-2 gene transcription in cells
20 derived from different species, providing a strategy for the study of gene function in a
variety of organisms.

EXAMPLE 11: Specific Induction of G1 accumulation of ErbB-2 overexpressing tumor
25 **cells.**

Overexpression of ErbB-2 leads to constitutive activation of its intrinsic tyrosine kinase
activity (P. P. Di Fiore *et al.*, *Science* 237, 178 (1987)), and it has been shown that
downregulation of ErbB-2 in tumor cells overexpressing the receptor leads to growth
30 inhibition (R. M. Hudziak *et al.*, *Mol. Cell. Biol.* 9, 1165 (1989); J. Deshane *et al.*, *Gene Ther.*
1, 332 (1994); J. M. Daly *et al.*, *Cancer Res.* 57, 3804 (1997)). The mechanism of growth
inhibition appears to be that progression of the cells from the G1 to the S phase of the cell
cycle is prevented (R. M. Neve, H. Sutterluty, N. Pullen, H. A. Lane, J. M. Daly, W. Krek, N.
35 E. Hynes, Submitted for publication). Thus, we investigated if expression of our designed
transcriptional repressor in erbB-2 overexpressing tumor cells would lead to a G1 block.
Therefore, SKBR3 breast cancer cells were infected with E2C-KRAB retrovirus and cell-
cycle distribution was analyzed in relation to ErbB-2 expression levels by flow cytometry
40 (22). Two cell populations were observed: about 40% of the cells were not infected and had
normal ErbB-2 levels, while the infected cells, ~60%, displayed approximately 7-fold
reduced receptor levels after 3 days. Compared to cells with normal receptor levels, a
significantly larger fraction of cells with decreased ErbB-2 expression levels was in the G1
45 phase of the cell cycle. To ascertain that the G1 accumulation observed with SKBR3 cells
was specific for ErbB-2 overexpressing tumor cells, a similar analysis was carried out with
the T47D breast cancer cell line, which does not display elevated levels of ErbB-2 (Fig. 4B).
50 Indeed, when T47D cells were infected with the E2C-KRAB retrovirus and subjected to flow

5

10

15

20

25

30

35

40

45

50

55

cytometric analysis, cell populations with normal and reduced ErbB-2 levels were found to display indistinguishable DNA content. Thus, our designed repressor protein is able to specifically induce G1 accumulation of ErbB-2 overexpressing tumor cells. The ability to inhibit cell-cycle progression, and hence inhibit growth of ErbB-2 overexpressing tumor cells suggests the potential of designed transcription factors for cancer gene therapy.

Claims

5

10

15

20

25

30

35

40

45

50

55

5

CLAIMS:

10

1. An isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of any of SEQ ID NO:1-16.

15

2. A composition comprising from 2 to about 12 of the polypeptide of claim 1.

3. The composition of claim 2 containing from 2 to about 6 polypeptides.

20

4. The composition of claim 2 or 3 wherein the polypeptides are operatively linked.

25

5. The composition of any of claims 2 to 4 wherein the polypeptides are linked by a linker having the sequence of SEQ ID NO 111.

30

6. The composition of any of claims 2 to 5 wherein each of the polypeptides binds to a different nucleotide sequence.

35

7. The composition of any of claims 2 to 6 that binds to a nucleotide that contains the sequence 5'-(GNN)_n-3', wherein each N is A, C, G, or T with the proviso that all N's cannot be C and where n is 2 to 6.

40

8. The polypeptide of claim 1 further operatively linked to one or more transcription regulating factors.

45

9. The composition of any of claims 2 to 7 further operatively linked to one or more transcription regulating factors.

50

10. An isolated and purified polynucleotide that encodes the polypeptide of claim 1.

11. An isolated and purified polynucleotide that encodes the composition of any of claims 2 to 7.

55

5

12. An expression vector containing the polynucleotide of claim 10.

10

13. An expression vector containing the polynucleotide of claim 11.

15

14. A process of regulating a nucleotide sequence that contains the sequence 5'-(GNN)_n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of the composition of any of claims 2 to 7.

20

15. The process of claim 14 wherein the sequence 5'-(GNN)_n-3' is located in the transcribed region of the nucleotide sequence.

25

16. The process of claim 14 wherein the sequence 5'-(GNN)_n-3' is located in a promotor region of the nucleotide sequence.

30

17. The process of claim 14 wherein the sequence 5'-(GNN)_n-3' is located within an expressed sequence tag.

35

18. The process of claim 14 wherein the composition is operatively linked to one or more transcription modulating factors.

40

19. Medicament comprising the composition of any of claims 2 to 7.

20. Use of . the composition of any of claims 2 to 7 for the manufacture of a medicament for the treatment of a human being.

45

21. Use of . the composition of any of claims 2 to 7 for the manufacture of a medicament for the treatment of cancer.

50

55

Fig. 1

Target 5'---3' (SEQ ID NO:)		Binding-helix amino acids at positions -1 1 2 3 4 5 6	Specificity
GAA	(1)	Q S S N L V R	GAA (GAT)
	(17)	Q R S N L V R	GAA, GAT
	(18)	Q S G N L V R	GAN
	(19)	Q P G N L V R	GAN
GAC	(2)	D P G N L V R	GAC
	(20)	D P G N L K R	GAC, GAT
GAG	(3)	R S D N L V R	GAG
	(21)	R S D N L R R	GAG, GGG
	(22)	K S A N L V R	GAG, (GAT)
	(23)	R S D N L V K	GAG, (GGG)
	(24)	K S A Q L V R	UNSPEC.
GAT	(4)	T S G N L V R	GAT
GCA	(5)	Q S G D L R R	GCA, GCT
	(25)	Q S S T L V R	GTA, GCA
	(26)	Q S G T L R R	GTA, GCA/T/C
	(27)	Q P G D L V R	GCT, GCC, GCA

Fig. 1

	(28)	Q G P D L V R	GCT, GCA
	(29)	Q A G T L M R	GTA, GCA
	(30)	Q P G T L V R	GTA, GCA
	(31)	Q G P E L V R	non-binder
GCC	(6)	D C R D L A R	GCC
	(32)	G C R E L S R	GCC
	(33)	D P S T L K R	GCC (GCA/T GTC)
	(34)	D P S D L K R	GCC, GAC
	(35)	D S G D L V R	GCC, GAC
	(36)	D S G E L V R	GCT, GCC
	(37)	D S G E L K R	GCT, GCC, GTC
GCG	(7)	R S D D L V K	GCG
	(38)	R L D T L G R	GNG
	(39)	R P G D L V R	GCG, GNG, GCN
	(40)	R S D T L V R	NG
	(41)	K S A D L K R	GAG, GTG, GCT, GCC
	(42)	R S D D L V R	GAG, (GNG, GCN)
	(43)	R S D T L V K	GNG

Fig. 1

	(44)	K S A E L K R	GCT, GCC, UNSPEC.
	(45)	K S A E L V R	GCT, GCC, UNSPEC.
	(46)	R G P E L V R	UNSPEC.
	(47)	K P G E L V R	NON-BINDER, BUT EXPR.
GCT	(8)	T S G E L V R	GCT
	(48)	S S Q T L T R	GCT
	(49)	T P G E L V R	GCT
	(50)	T S G D L V R	GCT, (GCC, GCA)
	(51)	S S Q T L V R	GCT
	(52)	T S Q T L T R	GCT (GAT, GTC, GCC)
	(53)	T S G E L K R	GCT, GCC
	(54)	Q S S D L V R	GCT (GCA, GCC)
	(55)	S S G T L V R	GCC, GCT
	(56)	T P G T L V R	GCT, GTC
	(57)	T S Q D L K R	GCC, GCT
	(58)	T S G T L V R	GCT, UNSPEC.
GGA	(9)	Q R A H L E R	GGA
	(59)	Q S S H L V R	GGA
	(60)	Q S G H L V R	GGA

Fig. 1

	(61)	Q P G H L V R	GGA, GCT
GGC	(10)	D P G H L V R	GGC
	(62)	E R S K L A R	GGC
	(63)	D P G H L A R	GGC
	(64)	Q R A K L E R	GGC
	(65)	Q S S K L V R	GGC
	(66)	D R S K L A R	GGC, GGN
	(67)	D P G K L A R	GGC, unspec.
GGG	(11)	R S D K L V R	GGG
	(68)	R S D K L T R	GGG
	(69)	R S D H L T R	GGG, GAG
	(70)	K S A K L E R	NON-BINDER
GGT	(12)	T S G H L V R	GGT, GGA
	(71)	T A D H L S R	GGT, GAT
	(72)	T A D K L S R	GGG, (GGT)
	(73)	T P G H L V R	GGT, unspec.
	(74)	T S S H L V R	unspec.
	(75)	T S G K L V R	unspec.
GTA	(13)	Q S S S L V R	
	(76)	Q P G E L V R	GTA, (GCT)
	(77)	Q S G E L V R	GTA, GCA/C

Fig. 1

	(78)	Q S G E L R R	GTA, GCA/T/C
GTC	(14)	D P G A L V R	
	(79)	D P G S L V R	GTC (GCT, GCC)
GTG	(15)	R S D E L V R	GTG, (GAG, GCG)
	(80)	R K D S L V R	GTG, GNG
	(81)	R S D V L V R	GTG, GAG, GGG
	(82)	R H D S L L R	GTG, GAG, GNG
	(83)	R S D A L V R	GAG, GTG, GGG
	(84)	R S S S L V R	GTG
	(85)	R S S S H V R	GTG, GGG
	(86)	R S D E L V K	GTG
	(87)	R S D A L V K	GAG GTG GGG
	(88)	R S D V L V K	GAG GNG
	(89)	R S S A L V R	GNG
	(90)	R K D S L V K	GGG GNG
	(91)	R S A S L V R	GAG, unspec.
	(92)	R S D S L V R	GCT unspec.
	(93)	R I H S L V R	unspec.

Fig. 1

	(94)	R P G S L V R	UNSPEC.
	(95)	R G P S L V R	UNSPEC.
	(96)	R P G A L V R	UNSPEC.
	(97)	K S A S L V R	NON-BINDER
	(98)	K S A A L V R	NON-BINDER
	(99)	K S A V L V R	NON-BINDER
GTT	(16)	T S G S L V R	GTT, GCT
	(100)	T S G S L T R	GGT, GCT
	(101)	T S Q S L V R	GAT, GTA GCT, GCA
	(102)	T S S S L V R	GTA, GAT
	(103)	T P G S L V R	GTA
	(104)	T S G A L V R	GGT, GCT, GAT
	(105)	T P G A L V R	GGT, GAT, GCT
	(106)	T G G S L V R	GGT, GAT
	(107)	T S G E L V R	GCT GCG GTA GTT
	(108)	T S G E L T R	GCT GTA/T/C
	(109)	T S S A L V K	UNSPEC
	(110)	T S S A L V R	UNSPEC



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/10, C07K 14/47	A3	(11) International Publication Number: WO 00/23464 (43) International Publication Date: 27 April 2000 (27.04.2000)
(21) International Application Number: PCT/EP99/07742 (22) International Filing Date: 14 October 1999 (14.10.1999) (30) Priority Data: 09/173,941 16 October 1998 (16.10.1998) US (60) Parent Application or Grant NOVARTIS AG [/]; (). NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [/]; (). THE SCRIPPS RESEARCH INSTITUTE [/]; (). BARBAS, Carlos, F. [/]; (). BARBAS, Carlos, F. [/]; (). BECKER, Konrad; ().	Published	
(54) Title: ZINC FINGER BINDING DOMAINS FOR GNN (54) Titre: DOMAINES DE LIAISON DES DOIGTS DE ZINC POUR GNN (57) Abstract Zinc finger-nucleotide binding polypeptides having binding specificity for target nucleotides containing one or GNN triplets are provided. Compositions containing such polypeptides and the use of such polypeptides and compositions for regulating nucleotide function are also provided. (57) Abrégé L'invention porte sur des polypeptides de liaison des doigts de zinc ayant une spécificité de liaison pour des nucléotides cibles contenant un ou des triplets de GNN. L'invention porte également sur des compositions contenant ces polypeptides et sur l'utilisation de ces polypeptides et de ces compositions pour réguler la fonction nucléotidique.		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/10, C07K 14/47	A3	(11) International Publication Number: WO 00/23464 (43) International Publication Date: 27 April 2000 (27.04.00)
(21) International Application Number: PCT/EP99/07742 (22) International Filing Date: 14 October 1999 (14.10.99) (30) Priority Data: 09/173,941 16 October 1998 (16.10.98) US (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (71) Applicant (for all designated States except US): THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, La Jolla, CA 92037 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): BARBAS, Carlos, F. [US/US]; 755 Pacific Surf Drive, Solana Beach, CA 92075 (US).	(74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HK, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. (88) Date of publication of the international search report: 16 November 2000 (16.11.00)	
(54) Title: ZINC FINGER BINDING DOMAINS FOR GNN (57) Abstract <p>Zinc finger-nucleotide binding polypeptides having binding specificity for target nucleotides containing one or GNN triplets are provided. Compositions containing such polypeptides and the use of such polypeptides and compositions for regulating nucleotide function are also provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/EP 99/07742A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/10 C07K14/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	T. OGAWA ET AL.: "Enhanced expression in seminoma of human zinc finger genes located on chromosome 19" CANCER GENET. CYTOGENET., vol. 100, no. 1, 1 January 1998 (1998-01-01), pages 36-42, XP000882080 * page 37, right column, last full-paragraph; figure 1A * --- -/--	1-11, 19-21

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

27 March 2000

Date of mailing of the international search report

23.06.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl
Fax: (+31-70) 340-3018

Authorized officer

Julia, P

INTERNATIONAL SEARCH REPORT

 Intern. Application No
 PCT/EP 99/07742

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
Y	B. GEBELEIN ET AL.,: "A novel profile of expressed tags for zinc finger encoding genes from the poorly differentiated exocrine pancreatic cell line AR4IP" CANCER LETT., vol. 105, no. 2, 2 August 1996 (1996-08-02), pages 225-231, XP000882088 * whole document, in particular Table 1 * ---	1-11, 19-21
A	CHOO Y ET AL: "Selection of DNA binding sites for zinc fingers using rationally randomized DNA reveals coded interactions" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA,US,NATIONAL ACADEMY OF SCIENCE. WASHINGTON, 8 November 1994 (1994-11-08), pages 1168-1172, XP002075339 ISSN: 0027-8424 cited in the application * whole document, in particular figure 1 and page 11172, left-column, last paragraph to right-column, last paragraph * ---	1-21
A	CHOO Y ET AL: "Toward a code for the interactions of zinc fingers with DNA: Selection of randomized fingers displayed on phage" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA,US,NATIONAL ACADEMY OF SCIENCE. WASHINGTON, 8 November 1994 (1994-11-08), pages 1163-1167, XP002075340 ISSN: 0027-8424 cited in the application * whole document, in particular figure 2 * ---	1-21
Y	WO 96 06166 A (MEDICAL RES COUNCIL ;CHOO YEN (SG); KLUG AARON (GB); GARCIA ISIDRO) 29 February 1996 (1996-02-29) * whole document, in particular Table 1 and figures 4 and 7 * ---	1-21
Y	Q. LIU ET AL.,: "Design of polydactyl zinc-finger proteins for unique addressing within complex genomes" PROC. NATL. ACAD. SCI. USA, vol. 94, May 1997 (1997-05), pages 5525-5530, XP002918175 cited in the application * whole document, in particular page 5529, right-column * -----	1-21

INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/EP 99/07742

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1 - 21 (partial)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 1; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 2-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 1; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

2. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 2; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1 or 3-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 2; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

3. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 3; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-2 or 4-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 3; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

compositions; medicament comprising these compositions and uses thereof

4. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 4; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-3 or 5-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 4; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

5. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 5; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-4 or 6-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 5; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

6. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 6; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-5 or 7-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 6; an isolated and purified

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

7. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 7; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-6 or 8-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 7; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

8. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 8; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-7 or 9-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 8; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

9. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 9; compositions comprising from 2 to about 12 of isolated and purified zinc

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-8 or 10-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 9; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

10. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 10; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-9 or 11-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 10; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

11. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 11; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-10 or 12-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 11; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

12. Claims: 1-21 (partial)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 12; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-11 or 13-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 12; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

13. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 13; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-12 or 14-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 13; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

14. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 14; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-13 or 15-16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 14; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

compositions; medicament comprising these compositions and uses thereof

15. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 15; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-14 or 16 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 15; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

16. Claims: 1-21 (partial)

an isolated and purified zinc finger-nucleotide binding polypeptide that contains a nucleotide binding region having the sequence of SEQ ID No.: 16; compositions comprising from 2 to about 12 of isolated and purified zinc finger-nucleotide binding polypeptides containing the nucleotide binding regions having the sequence of any of SEQ ID No.: 1-15 and wherein at least one of said polypeptides contains the nucleotide binding region having the sequence of SEQ ID No.: 16; an isolated and purified polynucleotide encoding said polypeptide or compositions thereof; expression vectors; a process of regulating a nucleotide sequence that contains 5'-(GNN)n-3', where n is an integer from 1 to 6, the process comprising exposing the nucleotide sequence to an effective amount of these compositions; medicament comprising these compositions and uses thereof

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/JP 99/07742

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9606166 A	29-02-1996	AU 1003799 A	22-04-1999
		AU 698152 B	22-10-1998
		AU 3229195 A	14-03-1996
		CA 2196419 A	29-02-1996
		EP 0781331 A	02-07-1997
		JP 10504461 T	06-05-1998
		US 6013453 A	11-01-2000
		US 6007988 A	28-12-1999
